

EULERIAN NUMBERS AND OPERATORS

by

L. CARLITZ

1. INTRODUCTION. The Eulerian numbers $A_{n,k}$ are usually introduced by means of [1], [6, Ch. 8]

$$(1.1) \quad \frac{1 - \lambda}{1 - \lambda e^{(1-\lambda)x}} = 1 + \sum_{n=1}^{\infty} \frac{x^n}{n!} \sum_{k=1}^n A_{n,k} x^k.$$

It follows from (1.1) that

$$(1.2) \quad A_{n+1,k} = (n - k + 2) A_{n,k-1} + k A_{n,k}$$

and

$$(1.3) \quad A_{n,k} = A_{n,n-k+1} \quad (1 \leq k \leq n).$$

It is evident from (1.2) and $A_{1,1} = 1$ that the $A_{n,k}$ are positive integers for $n \geq k \geq 1$.

The symmetry relation (1.3) is by no means obvious from the generating function (1.1). This has motivated the introduction of the symmetric notation [3]

$$(1.4) \quad A(r, s) = A_{r+s+1, r+1} = A_{r+s+1, s+1} = A(s, r),$$

where now $r \geq 0, s \geq 0$. It then follows from (1.1) that

$$(1.5) \quad \sum_{r,s=0}^{\infty} A(r, s) \frac{x^r y^s}{(r+s+1)!} = F(x, y),$$

where

$$(1.6) \quad F(x, y) = \frac{e^x - e^y}{x e^y - y e^x}$$

The recurrence (1.2) becomes

$$(1.7) \quad A(r, s) = (r+1)A(r, s-1) + (s+1)A(r-1, s).$$

Moreover in addition to (1.5) there is a second generating function

$$(1.8) \quad \sum_{r,s=0}^{\infty} A(r, s) \frac{x^r y^s}{(r+s)!} = (1 + xF(x, y))(1 + yF(x, y))$$

with $F(x, y)$ defined by (1.6).

If we put

$$(1.9) \quad A_n = A_n(x, y) = \sum_{r+s=n} A(r, s) x^r y^s,$$

it follows from (1.7) that

$$(1.10) \quad A_n(x, y) = (x + y + xy(D_x + D_y))A_{n-1}(x, y),$$

where $D_x = \partial/\partial x$, $D_y = \partial/\partial y$. Iteration of (1.10) gives

$$(1.11) \quad A_n(x, y) = (x + y + xy(D_x + D_y))^n \cdot 1.$$

It is accordingly of interest to consider the expansion of the operator

$$(1.12) \quad \Omega^n \equiv (x + y + xy(D_x + D_y))^n.$$

We shall show that

$$(1.13) \quad \Omega^n = \sum_{k=0}^n C_{n,k}(x, y) (xy)^k (D_x + D_y)^k,$$

where

$$(1.14) \quad C_{n,k}(x, y) = \frac{1}{k!(k+1)!} (D_x + D_y)^k A_n(x, y) \quad (0 \leq k \leq n).$$

The generating function (1.8) suggests the generalization [3]

$$(1.15) \quad \sum_{r,s=0}^{\infty} A(r, s | \alpha, \beta) \frac{x^r y^s}{(r+s)!} = (1 + xF(x, y))^\alpha (1 + yF(x, y))^\beta$$

where again $F(x, y)$ is defined by (1.6). Thus

$$A(r, s) = A(r, s|1, 1).$$

It follows from (1.15) that

$$(1.16) \quad A(r, s|\alpha, \beta) = (r + \beta) A(r, s - 1|\alpha, \beta) + (s + \alpha) A(r - 1, s|\alpha, \beta),$$

which evidently reduces to (1.7) when $\alpha = \beta = 1$; also

$$(1.17) \quad A(r, s|\alpha, \beta) = A(s, r|\beta, \alpha).$$

By (1.16), $A(r, s|\alpha, \beta)$ is a polynomial in α, β with positive integral coefficients. Combinatorial properties of $A(r, s|\alpha, \beta)$ are discussed in [3].

Put

$$(1.18) \quad A_n(x, y|\alpha, \beta) = \sum_{r+s=n} A(r, s|\alpha, \beta) x^r y^s.$$

Then by (1.16)

$$(1.19) \quad A_n(x, y|\alpha, \beta) = (\alpha x + \beta y + xy(D_x + D_y)) A_{n-1}(x, y|\alpha, \beta),$$

so that

$$(1.20) \quad A_n(x, y|\alpha, \beta) = (\alpha x + \beta y + xy(D_x + D_y))^n \cdot 1.$$

It is therefore of interest to consider the expansion of the operator

$$(1.21) \quad \Omega_{\alpha, \beta}^n \equiv (\alpha x + \beta y + xy(D_x + D_y))^n.$$

We shall show that

$$(1.22) \quad \Omega_{\alpha, \beta}^n = \sum_{k=0}^n C_{n, k}^{(\alpha, \beta)}(x, y) (xy)^k (D_x + D_y)^k,$$

where

$$(1.23) \quad C_{n, k}^{(\alpha, \beta)}(x, y) = \frac{1}{k! (\alpha + \beta)_k} (D_x + D_y)^k A_n(x, y|\alpha, \beta) \quad (0 \leq k \leq n),$$

where

$$(\alpha + \beta)_k = (\alpha + \beta)(\alpha + \beta + 1) \dots (\alpha + \beta + k - 1).$$

The case $\alpha + \beta$ equal to zero or a negative integer requires special treatment.

We consider also the inverse of (1.22), that is,

$$(1.24) \quad (xy)^n (D_x + D_y)^n = \sum_{k=0}^n B_{n,k}^{(\alpha,\beta)} \Omega_{\alpha,\beta}^k.$$

We show that

$$(1.25) \quad (D_x + D_y) B_{n,k}^{(\alpha,\beta)}(x, y) = n(\alpha + \beta + n - 1) B_{n-1,k}^{(\alpha,\beta)}(x, y)$$

and

$$(1.26) \quad \sum_{n=0}^{\infty} \frac{u^n}{n!} \sum_{k=0}^n B_{n,k}^{(\alpha,\beta)}(x, y) (x-y)^k v^k = \left(\frac{1-xu}{1-yu} \right)^{-v} (1-xu)^{-\alpha} (1-yu)^{-\beta}.$$

Additional properties of $B_{n,k}^{(\alpha,\beta)}(x, y)$ are given in §§ 8-10.

In recent years the Eulerian numbers and certain generalizations have been encountered in a number of combinatorial problems [2], [3], [4], [5], [6], [7]. The study of Eulerian operators is of intrinsic interest and may be useful for applications.

2. It is convenient to first discuss (1.13), that is,

$$(2.1) \quad (x + y + xy(D_x + D_y))^n = \sum_{k=0}^n C_{n,k} (x, y) (xy)^k (D_x + D_y)^k.$$

We shall require the following operational formulas:

$$(2.2) \quad (D_x + D_y)^k (x + y) = 2k(D_x + D_y)^{k-1} + (x + y)(D_x + D_y)^k,$$

$$(2.3) \quad (D_x + D_y)^k xy = k(k-1)(D_x + D_y)^{k-2} + k(x+y)(D_x + D_y)^{k-1} + xy(D_x + D_y)^k.$$

The proof is by induction on k . For (2.2) we have

$$\begin{aligned} (D_x + D_y)^k (x + y) &= 2k(D_x + D_y)^{k-1} + (D_x + D_y)(x + y)(D_x + D_y)^{k-1} \\ &= 2k(D_x + D_y)^{k-1} + [2 + (x + y)(D_x + D_y)](D_x + D_y)^{k-1} \\ &= 2(k+1)(D_x + D_y)^{k-1} + (x + y)(D_x + D_y)^{k-1}. \end{aligned}$$

As for (2.3), we have

$$\begin{aligned}
 (D_x + D_y)^{k+1} xy &= k(k-1)(D_x + D_y)^{k-1} + \\
 &+ k(D_x + D_y)(x+y)(D_x + D_y)^{k-1} + (D_x + D_y)xy(D_x + D_y)^k \\
 &= k(k-1)(D_x + D_y)^{k-1} + k[2 + (x+y)(D_x + D_y)](D_x + D_y)^{k-1} \\
 &\quad + [x+y+xy(D_x + D_y)](D_x + D_y)^k \\
 &= k(k+1)(D_x + D_y)^{k-1} + (k+1)xy(D_x + D_y)^k + xy(D_x + D_y)^{k+1}.
 \end{aligned}$$

Incidentally, the special case $k = 1$ of (2.3) may be noted:

$$(2.4) \quad (D_x + D_y)xy = x + y + xy(D_x + D_y) \equiv \Omega$$

Thus

$$(2.5) \quad \Omega^n = [(D_x + D_y)xy]^n.$$

We now apply Ω to both sides of (2.1). Then

$$\Omega^{n+1} = \sum_{k=0}^n \Omega \{C_{n,k}(x, y)(xy)^k\} (D_x + D_y)^k.$$

Since

$$\begin{aligned}
 &(D_x + D_y) \{C_{n,k}(x, y)(xy)^k\} \\
 &= k(xy)^{k-1}(x+y)C_{n,k}(x, y) \\
 &+ (xy)^k(D_x + D_y)C_{n,k}(x, y) + (xy)^kC_{n,k}(x, y)(D_y + D_x),
 \end{aligned}$$

it follows that

$$\begin{aligned}
 \Omega^{n+1} &= (x+y) \sum_{k=0}^{\infty} C_{n,k}(x, y)(xy)^k (D_x + D_y)^k \\
 &+ xy \sum_{k=0}^n \{k(xy)^{k-1}(x+y)C_{n,k}(x, y) + (xy)^k(D_x + D_y)C_{n,k}(x, y) \\
 &\quad + (xy)^kC_{n,k}(x, y)(D_x + D_y)\} (D_x + D_y)^k \\
 &= \sum_k (xy)^k \{[(k+1)(x+y) + xy(D_x + D_y)]C_{n,k}(x, y) \\
 &\quad + C_{n,k-1}(x, y)\} (D_x + D_y)^k
 \end{aligned}$$

We therefore have the recurrence

(2.6)

$$C_{n+1,k}(x, y) = [(k+1)(x+y) + xy(D_x + D_y)]C_{n,k}(x, y) + C_{n,k-1}(x, y).$$

This establishes the existence of the expansion (2.1) and indeed shows that $C_{n,k}(x, y)$ is a homogeneous polynomial in x, y of degree $n - k$.

In the next place we apply Ω to both sides of (6.1) but now on the right. Then

$$\Omega^{n+1} = \sum_{k=0}^n C_{n,k}(x, y) (xy)^k (D_x + D_y)^k [x + y + xy(D_x + D_y)].$$

Applying (2.2) and (2.3), we get

$$\begin{aligned} \Omega^{n+1} &= \sum_{k=0}^n C_{n,k}(x, y) (xy)^k \{2k(D_x + D_y)^{k-1} + (x + y)(D_x + D_y)^k\} \\ &+ \sum_{k=0}^n C_{n,k}(x, y) (xy)^k \{k(k-1)(D_x + D_y)^{k-2} + k(x + y)(D_x + D_y)^{k-1} \\ &\quad + xy(D_x + D_y)^k\} (D_x + D_y). \end{aligned}$$

It follows that

$$(2.7) \quad \begin{aligned} C_{n+1,k}(x, y) &= (k+1)(x+y)C_{n,k}(x, y) \\ &+ (k+1)(k+2)xyC_{n,k+1}(x, y) + C_{n,k-1}(x, y). \end{aligned}$$

Comparing (2.7) with (2.6), we get

$$(2.8) \quad (D_x + D_y)C_{n,k}(x, y) = (k+1)(k+2)C_{n,k+1}(x, y).$$

It is clear from (2.8) that

$$(2.9) \quad C_{n,k}(x, y) = \frac{1}{k!(k+1)!} (D_x + D_y)^k C_{n,0}(x, y).$$

Since, by (1.1),

$$C_{n,0}(x, y) = A_n(x, y),$$

(2.9) becomes

$$(2.10) \quad C_{n,k}(x, y) = \frac{1}{k!(k+1)!} (D_x + D_y)^k A_n(x, y) \quad (0 \leq k \leq n).$$

so that we have proved (1.14).

3. Put

$$(3.1) \quad f_n(x, y, z) = \sum_{k=0}^n (k+1)! C_{n,k}(x, y) z^k.$$

Then $f_n(x, y, z)$ is homogeneous in x, y, z of degree n . We also define

$$(3.2) \quad g_k(x, y) = \sum_{n=k}^{\infty} \frac{1}{(n+1)!} C_{n,k}(x, y).$$

Since, by (1.5) and (1.9),

$$\sum_{n=0}^{\infty} \frac{1}{(n+1)!} A_n(x, y) = F(x, y),$$

it follows that

$$(3.3) \quad g_k(x, y) = \frac{1}{k!(k+1)!} (D_x + D_y)^k F(x, y).$$

It is easily verified that

$$(D_x + D_y) F = F^2$$

and therefore

$$(3.4) \quad (D_x + D_y)^k F = k! F^{k+1}.$$

Thus (3.3) becomes

$$(3.5) \quad g_k(x, y) = \frac{1}{(k+1)!} F^{k+1}(x, y).$$

Therefore

$$(3.6) \quad G(x, y, z) \equiv \sum_{k=0}^{\infty} (k+1)! g_k(x, y) z = \frac{F(x, y)}{1 - z F(x, y)}.$$

Also, since

$$G(x, y, z) = \sum_{n=0}^{\infty} \frac{1}{(n+1)!} f_n(x, y, z),$$

we get

$$(3.7) \quad \sum_{n=0}^{\infty} \frac{1}{(n+1)!} f_n(x, y, z) = \frac{F(x, y)}{1 - z F(x, y)}.$$

By (1.6),

$$\begin{aligned} \frac{F(x, y)}{1 - z F(x, y)} &= \frac{e^x - e^y}{(xe^y - ye^x) - z(e^x - e^y)} \\ &= \frac{e^x - e^y}{(x+z)e^x - (y+z)e^y} \\ &= \frac{e^{x+z} - e^{y+z}}{(x+z)e^{y+z} - (y+z)e^{x+z}} = F(x+z, y+z). \end{aligned}$$

Thus (3.7) becomes

$$(3.8) \quad \sum_{n=0}^{\infty} \frac{1}{(n+1)!} f_n(x, y, z) = F(x+z, y+z).$$

Since

$$F(x+z, y+z) = \sum_{n=0}^{\infty} A_n(x+z, y+z),$$

it follows that

$$(3.9) \quad f_n(x, y, z) = A_n(x+z, y+z).$$

This formula can also be proved without the use of generating functions.

4. We now consider the general case:

$$(4.1) \quad \Omega_{\alpha, \beta}^n = \sum_{k=0}^n C_{n, k}^{(\alpha, \beta)}(x, y) (xy)^k (D_x + D_y)^k,$$

where

$$(4.2) \quad \Omega_{\alpha, \beta} \equiv \alpha x + \beta y + xy(D_x + D_y).$$

We apply $\Omega_{\alpha, \beta}$ on the left of each side of (4.1). Since

$$\begin{aligned} & \Omega_{\alpha, \beta} \{ C_{n, k}^{(\alpha, \beta)}(x, y) (xy)^k (D_x + D_y)^k \} \\ &= (\alpha x + \beta y) C_{n, k}^{(\alpha, \beta)}(x, y) (xy)^k + k(x+y) C_{n, k}^{(\alpha, \beta)}(x, y) (xy)^k \\ &+ (xy)^{k+1} (D_x + D_y) C_{n, k}^{(\alpha, \beta)}(x, y) + (xy)^{k+1} C_{n, k}^{(\alpha, \beta)}(x, y) (D_x + D_y), \end{aligned}$$

we get the recurrence

$$(4.3) \quad C_{n+1, k}^{(\alpha, \beta)}(x, y) = (\alpha x + \beta y) C_{n, k}^{(\alpha, \beta)}(x, y) \\ + [k(x+y) + xy(D_x + D_y)] C_{n, k}^{(\alpha, \beta)}(x, y) + C_{n, k-1}^{(\alpha, \beta)}(x, y).$$

Next, apply $\Omega_{\alpha, \beta}$ on the right. Since

$$\begin{aligned} (D_x + D_y)^k (\alpha x + \beta y) &= \sum_{j=0}^k \binom{k}{j} D_x^j D_y^{k-j} (\alpha x + \beta y) \\ &= \sum_{j=0}^k \binom{k}{j} \{ \alpha (x D_x^j + j D_x^{j-1}) D_y^{k-j} + \beta D_x^j (y D_y^{k-j} + (k-j) D_y^{k-j-1}) \} \end{aligned}$$

$$\begin{aligned}
&= \alpha x \sum_{j=0}^k \binom{k}{j} D_x^j D_y^{k-j} + k \alpha \sum_{j=1}^k \binom{k-1}{j-1} D_x^{j-1} D_y^{k-j} + \beta y \sum_{j=0}^k \binom{k}{j} D_x^j D_y^{k-j} \\
&\quad + k \beta \sum_{j=0}^{k-1} \binom{k-1}{j} D_x^j D_y^{k-j} \\
&= (\alpha x + \beta y) (D_x + D_y)^k + k(\alpha + \beta) (D_x + D_y)^{k-1}
\end{aligned}$$

and, by (2.3),

$$\begin{aligned}
&(D_x + D_y)^k xy + k(k-1) (D_x + D_y)^{k-2} \\
&+ k(x+y) (D_x + D_y)^{k-1} + xy (D_x + D_y)^k,
\end{aligned}$$

we get

$$\begin{aligned}
\Omega_{\alpha, \beta}^{n+1} &= \sum_{k=0}^n C_{n, k}^{(\alpha, \beta)}(x, y) (xy)^k \{(\alpha x + \beta y) (D_x + D_y)^k \\
&\quad + k(\alpha + \beta) + (D_x + D_y)^{k-1} \\
&+ k(k-1) (D_x + D_y)^{k-1} + k(x+y) (D_x + D_y)^k + xy (D_x + D_y)^{k+1}\}.
\end{aligned}$$

It follows that

$$\begin{aligned}
(4.4) \quad C_{n+1, k}^{(\alpha, \beta)}(x, y) &= ((k + \alpha)x + (k + \beta)y) C_{n, k}^{(\alpha, \beta)}(x, y) \\
&+ (k + 1)(k + \alpha + \beta) xy C_{n, k+1}^{(\alpha, \beta)}(x, y) + C_{n, k-1}^{(\alpha, \beta)}(x, y).
\end{aligned}$$

Comparison of (4.4) with (4.3) gives

$$(4.5) \quad (D_x + D_y) C_{n, k}^{(\alpha, \beta)}(x, y) = (k + 1)(k + \alpha + \beta) C_{n, k-1}^{(\alpha, \beta)}(x, y).$$

It follows that

$$(4.6) \quad C_{n, k}^{(\alpha, \beta)}(x, y) = \frac{1}{k! (\alpha + \beta)_k} (D_x + D_y)^k C_{n, 0}^{(\alpha, \beta)}(x, y),$$

provided $\alpha + \beta$ is not equal to zero or a negative integer. Moreover, by (1.20),

$$C_{n, 0}^{(\alpha, \beta)}(x, y) = A_n(x, y | \alpha, \beta),$$

so that (4.6) becomes

$$(4.7) \quad C_{n, k}^{(\alpha, \beta)}(x, y) = \frac{1}{k! (\alpha + \beta)_k} (D_x + D_y)^k A_n(x, y | \alpha, \beta).$$

It follows from (1.19) and (4.7) that

$$(4.8) \quad A_{m+n}(x, y | \alpha, \beta) \\ = \sum_{k=0}^{\min(m, n)} \frac{1}{k! (\alpha + \beta)_k} (xy)^k (D_x + D_y)^k A_m(x, y | \alpha, \beta) (D_x + D_y)^k A_n(x, y | \alpha, \beta).$$

5. Put

$$(5.1) \quad f_n(x, y, z | \alpha, \beta) = \sum_{k=0}^n (\alpha + \beta)_k C_{n, k}^{(\alpha, \beta)}(x, y) z^k,$$

$$(5.2) \quad g_k(x, y | \alpha, \beta) = \sum_{n=k}^{\infty} \frac{1}{n!} C_{n, k}^{(\alpha, \beta)}(x, y),$$

$$(5.3) \quad \Phi_{\alpha, \beta}(x, y) = (1 + xF(x, y))^\alpha (1 + yF(x, y))^\beta.$$

Since

$$\sum_{n=0}^{\infty} \frac{1}{n!} A_n(x, y | \alpha, \beta) = \Phi_{\alpha, \beta}(x, y),$$

it follows from (4.7) and (5.2) that

$$(5.4) \quad g_k(x, y) = \frac{1}{k! (\alpha + \beta)_k} (D_x + D_y)^k \Phi_{\alpha, \beta}(x, y).$$

But

$$(5.5) \quad (D_x + D_y)^k \Phi_{\alpha, \beta}(x, y) = (\alpha + \beta)_k F^k(x, y) \Phi_{\alpha, \beta}(x, y),$$

so that (5.4) becomes

$$(5.6) \quad g_k(x, y | \alpha, \beta) = \frac{1}{k!} F^k(x, y) \Phi_{\alpha, \beta}(x, y).$$

Now put

$$G(x, y, z | \alpha, \beta) = \sum_{k=0}^{\infty} (\alpha + \beta)_k g_k(x, y | \alpha, \beta) z^k.$$

Then, by (5.6),

$$(5.7) \quad G(x, y, z | \alpha, \beta) = \frac{\Phi_{\alpha, \beta}(x, y)}{(1 - zF(x, y, y))^{\alpha + \beta}}.$$

Since

$$\Phi_{\alpha, \beta}(x, y) = \frac{(x - y)^{\alpha + \beta} e^{x+y}}{(x e^x - y e^y)^{\alpha + \beta}}$$

and

$$\begin{aligned} \frac{\Phi_{\alpha,\beta}(x,y)}{(1-zF(x,y))^{\alpha+\beta}} &= \frac{(x-y)^{\alpha+\beta} e^{\alpha x + \beta y}}{[(xe^y - ye^x) - z(e^x - e^y)]} \\ &= \frac{(x-y)^{\alpha+\beta} e^{\alpha x + \beta y}}{[(x+z)e^y - (y+z)e^x]} \\ &= \frac{(x-y)^{\alpha+\beta} e^{\alpha(x+z) + \beta(x+z)}}{[(x+z)e^{y+z} - (y+z)e^{x+z}]^{\alpha+\beta}}, \end{aligned}$$

(5.7) becomes

$$(5.8) \quad G(x, y, z | \alpha, \beta) = \Phi_{\alpha,\beta}(x+z, y+z).$$

On the other hand, by (5.1) and (5.2),

$$\begin{aligned} G(x, y, z | \alpha, \beta) &= \sum_{k=0}^{\infty} (\alpha + \beta)_k z^k \sum_{n=k}^{\infty} \frac{1}{n!} C_{n,k}^{(\alpha,\beta)}(x, y) \\ &= \sum_{k=0}^{\infty} \frac{1}{k!} \sum_{n=k}^{\infty} (\alpha + \beta)_k C_{n,k}^{(\alpha,\beta)}(x, y) z^k \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} f_n(x, y, z | \alpha, \beta), \end{aligned}$$

so that, by (5.8),

$$(5.9) \quad \sum_{n=0}^{\infty} \frac{1}{n!} f_n(x, y, z | \alpha, \beta) = \Phi_{\alpha,\beta}(x+z, y+z).$$

Therefore, by (1.15) and (1.18), we get

$$(5.10) \quad f_n(x, y, z | \alpha, \beta) = A_n(x+z, y+z | \alpha, \beta).$$

This identity implies

$$(5.11) \quad A_n(x+z, y+z | \alpha, \beta) = \frac{z^k}{k!} (D_x + D_y)^k A_n(x, y | \alpha, \beta),$$

which can also be obtained by applying Taylor's theorem to $A_n(x, y | \alpha, \beta)$.

6. As noted above, (4.7) is not valid when $\alpha + \beta$ is zero or a negative integer. We shall now consider the excluded values. It is

convenient to begin with the special case $\alpha = \beta = 0$. In place of (4.1) we now have

$$(6.1) \quad (xy(D_x + D_y))^n = \sum_{k=1}^n C_{n,k}^{(0,0)}(x, y) (xy)^k (D_x + D_y) \quad (n \geq 1)$$

The recurrence (4.3) reduces to

$$(6.2) \quad C_{n+1,k}^{(0,0)}(x, y) = [k(x+y) + xy(D_x + D_y)] C_{n,k}^{(0,0)}(x, y) + C_{n,k-1}^{(0,0)}(x, y),$$

while (4.4) becomes

$$(6.3) \quad \begin{aligned} C_{n+1,k}^{(0,0)}(x, y) &= k(x+y) C_{n,k}^{(0,0)}(x, y) \\ &+ k(k+1)xy C_{n,k+1}^{(0,0)}(x, y) + C_{n,k-1}^{(0,0)}(x, y) \end{aligned}$$

Hence

$$(6.4) \quad (D_x + D_y) C_{n,k}^{(0,0)}(x, y) = k(k+1) C_{n,k+k}^{(0,0)}(x, y),$$

so that

$$(6.5) \quad C_{n,k}^{(0,0)}(x, y) = \frac{1}{k!(k-1)!} (D_x + D_y)^{k-1} C_{n,1}^{(0,0)}(x, y) \quad (k \geq 1)$$

For $k = 1$, (6.2) reduces to

$$C_{n+1,1}^{(0,0)}(x, y) = [x + y + xy(D_x + D_y)] C_{n,1}^{(0,0)}(x, y),$$

which yields

$$C_{n,1}^{(0,0)}(x, y) = [x + y + xy(D_x + D_y)]^{n-1} \cdot C_{1,1}^{(0,0)}(x, y)$$

It is clear from (6.1) that $C_{1,1}^{(0,0)}(x, y) = 1$ and therefore

$$(6.6) \quad C_{n,1}^{(0,0)}(x, y) = A_{n-1}(x, y) \equiv A_{n-1}(x, y | 1, 1)$$

Thus (6.5) becomes

$$(6.7) \quad C_{n,k}^{(0,0)}(x, y) = \frac{1}{k!(k-1)!} (D_x + D_y)^{k-1} A_{n-1}(x, y)$$

Before discussing the general case $\alpha + \beta$ equal to zero or a negative integer, we consider the expansion

$$(6.8) \quad \Omega_{\alpha, \beta}^n = \sum_{k=0}^n Q_{n, k}^{(\alpha, \beta)}(x, y) (xy (D_x + D_y))^k,$$

where the $Q_{n, k}^{(\alpha, \beta)}(x, y)$ are to be determined. Clearly

$$\begin{aligned} \Omega_{\alpha, \beta}^{n+1} &= (\alpha x + \beta y) \sum_{k=0}^n Q_{n, k}^{(\alpha, \beta)}(x, y) (xy (D_x + D_y))^k \\ &\quad + xy \sum_{k=0}^n (D_x + D_y) Q_{n, k}^{(\alpha, \beta)}(x, y) \cdot (xy (D_x + D_y))^k \\ &\quad + \sum_{k=0}^n Q_{n, k}^{(\alpha, \beta)}(x, y) (xy (D_x + D_y))^k, \end{aligned}$$

so that

$$(6.9) \quad Q_{n+1, k}^{(\alpha, \beta)}(x, y) = \Omega_{\alpha, \beta} Q_{n, k}^{(\alpha, \beta)}(x, y) + Q_{n, k-1}^{(\alpha, \beta)}(x, y)$$

For $k = 0$, (6.9) reduces to

$$Q_{n+1, 0}^{(\alpha, \beta)}(x, y) = \Omega_{\alpha, \beta} Q_{n, 0}^{(\alpha, \beta)}(x, y)$$

Since, by (6.8),

$$Q_{1, 0}^{(\alpha, \beta)}(x, y) = \alpha x + \beta y = A_1(x, y | \alpha, \beta),$$

it is clear that

$$(6.10) \quad Q_{n, 0}^{(\alpha, \beta)}(x, y) = A_n(x, y | \alpha, \beta)$$

We shall now show that

$$(6.11) \quad Q_{n, 0}^{(\alpha, \beta)}(x, y) = \binom{n}{k} A_{n-k}(x, y | \alpha, \beta) \quad (0 \leq k \leq n)$$

Clearly (6.11) holds for $n = 0$. Assuming that it holds up to and including the value n , we have by (6.9),

$$\begin{aligned}
Q_{n+1,k}^{(\alpha,\beta)}(x,y) &= \binom{n}{k} \Omega_{\alpha,\beta} A_{n-k}(x,y|\alpha,\beta) + \binom{n}{k-1} A_{n-k+1}(x,y|\alpha,\beta) \\
&= \binom{n}{k} A_{n-k+1}(x,y|\alpha,\beta) + \binom{n}{k-1} A_{n-k+1}(x,y|\alpha,\beta) \\
&= \binom{n}{k} A_{n-k+1}(x,y|\alpha,\beta)
\end{aligned}$$

We have therefore proved

$$(6.12) \quad \Omega_{\alpha,\beta}^n = \sum_{k=0}^n \binom{n}{k} A_{n-k}(x,y|\alpha,\beta) (xy(D_x + D_y))^k$$

This suggests the following more general result:

$$(6.13) \quad \Omega_{\alpha+\gamma,\beta+\delta}^n = \sum_{k=0}^n \binom{n}{k} A_{n-k}(x,y|\alpha,\beta) \Omega_{\gamma,\delta}^k$$

To prove (6.13), consider

$$\Omega_{\alpha+\gamma,\beta+\delta}^n = \sum_{k=0}^n R_{n,k} \Omega_{\gamma,\delta}^k,$$

where the $R_{n,k}$ are functions of $x, y, \alpha, \beta, \gamma, \delta$. Then

$$\begin{aligned}
\Omega_{\alpha+\gamma,\beta+\delta}^{n+1} &= [(\alpha + \gamma)x + (\beta + \delta)y + xy(D_x + D_y)] \sum_{k=0}^n R_{n,k} \Omega_{\gamma,\delta}^k \\
&= [(\alpha + \gamma)x + (\beta + \delta)] \sum_{k=0}^n R_{n,k} \Omega_{\gamma,\delta}^k \\
&\quad + xy \sum_{k=0}^n (D_x + D_y) R_{n,k} \cdot \Omega_{\gamma,\delta}^k + \sum_{k=0}^n R_{n,k} \cdot xy(D_x + D_y) \Omega_{\gamma,\delta}^k \\
&= \sum_{k=0}^n \Omega_{\alpha,\beta} R_{n,k} \cdot \Omega_{\gamma,\delta}^k + \sum_{k=0}^n R_{n,k} \Omega_{\gamma,\delta}^k
\end{aligned}$$

This evidently implies

$$(6.14) \quad R_{n+1,k} = \Omega_{\alpha,\beta} R_{n,k} + R_{n,k-1}.$$

Then, exactly as above, we show first that

$$R_{n,0} = A_n(x,y|\alpha,\beta)$$

and generally

$$R_{n,k} = \binom{n}{k} A_{n-k}(x, y | \alpha, \beta) \quad (0 \leq k \leq n)$$

This completes the proof of (6.13)

As a special case of (6.13), we note

$$(6.15) \quad (xy(D_x + D_y))^n = \sum_{k=0}^n \binom{n}{k} A_{n-k}(x, y | -\alpha, -\beta) \Omega_{\alpha,\beta}^k.$$

7 We now treat the general excluded case in (4.7), $\alpha + \beta$ equal to zero or a negative integer. We shall require the following formulas:

$$(7.1) \quad \begin{aligned} \Phi_{\alpha,\beta}(x, y) &= \sum_0^{\infty} \frac{1}{n!} A_n(x, y | \alpha, \beta) \\ &= (1 + xF(x, y))^\alpha (1 + yF(x, y))^\beta, \end{aligned}$$

$$(7.2) \quad (D_x + D_y)^k F(x, y) = k! F^{k+1}(x, y),$$

$$(7.3) \quad (D_x + D_y)^k \Phi_{\alpha,\beta}(x, y) = (\alpha + \beta)_k F^k(x, y) \Phi_{\alpha,\beta}(x, y).$$

If $\alpha + \beta$ is not equal to zero or a negative integer, we have seen that

$$C_{n,k}^{(\alpha,\beta)}(x, y) = \frac{1}{k! (\alpha + \beta)_k} (D_x + D_y)^k A_n(x, y | \alpha, \beta).$$

It follows that

$$\sum_{n=k}^{\infty} \frac{1}{n!} C_{n,k}^{(\alpha,\beta)}(x, y) = \frac{1}{k! (\alpha + \beta)_k} (D_x + D_y)^k \Phi_{\alpha,\beta}(x, y)$$

and therefore, by (7.3),

$$(7.4) \quad \sum_{n=k}^{\infty} \frac{1}{n!} C_{n,k}^{(\alpha,\beta)}(x, y) = \frac{1}{k!} F^k(x, y) \Phi_{\alpha,\beta}(x, y).$$

Put

$$(7.5) \quad F^k(x, y) = \sum_{n=0}^{\infty} \frac{1}{(n+k)!} A_n^{(k)}(x, y) \quad (k = 1, 2, 3, \dots),$$

where $A_n^{(k)}(x, y)$ is homogeneous of degree n in x, y . For $k = 1$ we have

$$(7.6) \quad A_n^{(1)}(x, y) = A_n(x, y) = A_n(x, y | 1, 1).$$

It follows from (7.5) and (7.6) that

$$(7.7) \quad A_n^{(k+1)}(x, y) = \sum_{r=0}^n \binom{n+k+1}{r+1} A_r(x, y) A_{n-r}^{(k)}(x, y)$$

and therefore the coefficients in $A_n^{(k)}(x, y)$ are positive integers

By (7.4) and (7.5) we get, since $C_{n,k}^{(\alpha,\beta)}(x, y)$ is of degree $n - k$,

$$(7.8) \quad C_{n,k}^{(\alpha,\beta)}(x, y) = \frac{1}{k!} \sum_{r=k}^n \binom{n}{r} A_{r-k}^{(k)}(x, y) A_{n-r}(x, y | \alpha, \beta).$$

Since both $C_{n,k}^{(\alpha,\beta)}(x, y)$ and $A_n(x, y | \alpha, \beta)$ are polynomials in α, β (as well as in x, y), it follows that (7.8) is valid for all α, β . The numerical coefficients in $C_{n,k}^{(\alpha,\beta)}(x, y)$ are integers; however that is not obvious from (7.8)

Since

$$A_n(x, y | 0, 0) = 0 \quad (n > 0),$$

it is evident that (7.8) implies

$$(7.9) \quad C_{n,k}^{(0,0)}(x, y) = \frac{1}{k!} A_{r-k}^{(k)}(x, y).$$

By (7.2) and (7.5) we have

$$(k-1)! \sum_{n=0}^{\infty} \frac{1}{(n+k)!} A_n^{(k)}(x, y) = (D_x + D_y)^{k-1} \sum_{n=0}^{\infty} \frac{1}{(n+1)!} A_n(x, y),$$

so that

$$(7.10) \quad A_n^{(k)}(x, y) = \frac{1}{(k-1)!} (D_x + D_y)^{k-1} A_{n+k-1}(x, y) \quad (1 \leq k \leq n).$$

Thus (7.8) becomes

$$(7.11) \quad C_{n,k}^{(\alpha,\beta)}(x, y) = \frac{1}{k!(k-1)!} \sum_{r=k}^n \binom{n}{r} (D_x + D_y)^{k-1} A_{r-1}(x, y) \\ \cdot A_{n-r}(x, y | \alpha, \beta) \quad (1 \leq k \leq n).$$

For $\alpha = \beta = 0$, (7.11) reduces to

$$(7.12) \quad C_{n,k}^{(0,0)}(x, y) = \frac{1}{k!(k-1)!} (D_x + D_y)^{k-1} A_{n-1}(x, y)$$

in agreement with (6.7)

Both (7.8) and (7.11) are valid for all α, β .

8. We now consider the inverse of (4.1), that is,

$$(8.1) \quad (xy)^n (D_x + D_y)^n = \sum_{k=0}^n (-1)^{n-k} B_{n,k}^{(\alpha,\beta)}(x, y) \Omega_{\alpha,\beta}^k,$$

where, as will appear presently, $B_{n,k}^{(\alpha,\beta)}(x, y)$ is a homogeneous polynomial in x, y of degree $n - k$. The existence of a formula of this kind is evidently implied by (4.1). For example

$$\begin{aligned} xy(D_x + D_y) &= -(\alpha x + \beta y) + \Omega_{\alpha,\beta}, \\ (xy)^2(D_x + D_y)^2 &= (\alpha x + \beta y)^2 + \alpha x^2 + \beta y^2 \\ &\quad - [(2\alpha + 1)x + (2\beta + 1)y] \Omega_{\alpha,\beta} + \Omega_{\alpha,\beta}^2. \end{aligned}$$

To get a recurrence for the coefficients $B_{n,k}^{(\alpha,\beta)}$ we apply the operator $xy(D_x + D_y)$ to both sides of (8.1) — on the left. This gives

$$\begin{aligned} &n(x+y)(xy)^n(D_x + D_y) + (xy)^{n+1}(D_x + D_y)^{n+1} \\ &= xy \sum_{k=0}^n (-1)^{n-k} \{ (D_x + D_y) B_{n,k}^{(\alpha,\beta)}(x, y) + B_{n,k}^{(\alpha,\beta)}(x, y) (D_x + D_y) \} \Omega_{\alpha,\beta}^k \\ &= \sum_{k=0}^n (-1)^{n-k} \{ xy(D_x + D_y) B_{n,k}^{(\alpha,\beta)}(x, y) - (\alpha x + \beta y) B_{n,k}^{(\alpha,\beta)}(x, y), \\ &\quad + B_{n,k}^{(\alpha,\beta)}(x, y) \Omega_{\alpha,\beta} \} \Omega_{\alpha,\beta}^k. \end{aligned}$$

It follows that

$$\begin{aligned} (xy)^{n+1}(D_x + D_y)^{n+1} &= -n(x+y) \sum_{k=0}^n (-1)^{n-k} B_{n,k}^{(\alpha,\beta)}(x, y) \Omega_{\alpha,\beta}^k \\ &\quad + \sum_{k=0}^n (-1)^{n-k} \{ xy(D_x + D_y) B_{n,k}^{(\alpha,\beta)}(x, y) \\ &\quad - (\alpha x + \beta y) B_{n,k}^{(\alpha,\beta)}(x, y) \} \Omega_{\alpha,\beta}^k \\ &\quad + \sum_{k=1}^{n+1} (-1)^{n-k+1} B_{n,k-1}^{(\alpha,\beta)}(x, y) \Omega_{\alpha,\beta}^k. \end{aligned}$$

Therefore

$$(8.2) \quad B_{n+1,k}^{(\alpha,\beta)}(x,y) = [(\alpha+n)x + (\beta+n)y - xy(D_x + D_y)] \\ \cdot B_{n,k}^{(\alpha,\beta)}(x,y) + B_{n,k-1}^{(\alpha,\beta)}(x,y).$$

On the other hand, if we multiply both sides of (8.1) on the right by $\Omega_{\alpha,\beta}$ we get

$$\sum_{k=0}^n (-1)^{n-k} B_{0,k}^{(\alpha,\beta)}(x,y) \Omega_{\alpha,\beta}^{k+1} \\ = (xy)^n (D_x + D_y)^n [\alpha x + \beta y + xy(D_x + D_y)] \\ = (xy)^n \{n(\alpha + \beta)(D_x + D_y)^{n-1} + (\alpha x + \beta y)(D_x + D_x)^n\} \\ + (xy)^n \{n(n-1)(D_x + D_y)^{n-1} + n(x+y)(D_x + D_y)^n \\ + xy(D_x + D_y)^{n+1}\} \\ = n(\alpha + \beta + n - 1)(xy)^n (D_x + D_y)^{n-1} \\ + [(\alpha + n)x + (\beta + n)y](xy)^n (D_x + D_y)^n + (xy)^{n+1} (D_x + D_y)^{n+1}.$$

This implies

$$(8.3) \quad B_{n+1,k}^{(\alpha,\beta)}(x,y) = [(\alpha + n)x + (\beta + n)y] B_{n,k}^{(\alpha,\beta)}(x,y) \\ - n(\alpha + \beta + n - 1)xy B_{n-1,k}^{(\alpha,\beta)} + B_{n,k-1}^{(\alpha,\beta)}(x,y).$$

Comparing (8.3) with (8.2), we get

$$(8.4) \quad (D_x + D_y) B_{n,k}^{(\alpha,\beta)}(x,y) = n(\alpha + \beta + n - 1) B_{n-1,k}^{(\alpha,\beta)}(x,y).$$

In the next place, it follows at once from (4.1) and (8.1) that

$$(8.5) \quad \sum_{k=j}^n (-1)^{k-j} C_{n,k}^{(\alpha,\beta)}(x,y) B_{n,j}^{(\alpha,\beta)}(x,y) = \delta_{n,j}$$

and

$$(8.6) \quad \sum_{k=j}^n (-1)^{n-k} B_{n,k}^{(\alpha,\beta)}(x,y) C_{k,j}^{(\alpha,\beta)}(x,y) = \delta_{n,j}.$$

A formula of a different kind can be obtained by applying each side of (8.1) to $A_r(x,y|\alpha,\beta)$. Since

$$(D_x + D_y)^n A_r(x,y|\alpha,\beta) = 0 \quad (0 \leq r < n), \\ \Omega_{\alpha,\beta}^k A_r(x,y|\alpha,\beta) = A_{r+k}(x,y|\alpha,\beta),$$

we get

$$(8.7) \quad \sum_{k=0}^n (-1)^{n-k} B_{n,k}^{(\alpha,\beta)}(x, y) A_{r+k}(x, y | \alpha, \beta) = 0 \quad (0 \leq r < n).$$

In either (8.5) or (8.6) take $j = n$. Since $C_{n,n}^{(\alpha,\beta)}(x, y) = 1$, we have

$$(8.8) \quad B_{n,n}^{(\alpha,\beta)}(x, y) = 1.$$

Also it is easily verified that

$$\Omega_{\alpha,\beta}(x^{-\beta}y^{-\alpha}) = 0,$$

so that

$$\Omega_{\alpha,\beta}^k(x^{-\beta}y^{-\beta}) = 0 \quad (k = 1, 2, 3, \dots).$$

Thus (8.1) implies

$$(-1)^n B_{n,o}^{(\alpha,\beta)}(x, y) = (xy)^n (D_x + D_y)^n x^{-\beta}y^{-\alpha}.$$

A little manipulation leads to

$$(8.9) \quad B_{n,o}^{(\alpha,\beta)}(x, y) = \sum_{k=0}^n \binom{n}{k} (\alpha)_k (\beta)_{n-k} x^k y^{n-k}.$$

This is equivalent to

$$(8.10) \quad \sum_{n=0}^{\infty} B_{n,o}^{(\alpha,\beta)}(x, y) \frac{n!}{z^n} = (1 - xz)^{-\alpha} (1 - yz)^{-\beta}.$$

We remark that $B_{n,o}^{(\alpha,\beta)}(x, y)$ satisfies the following recurrence:

$$(8.11) \quad B_{n+1,o}^{(\alpha,\beta)}(x, y) = (\alpha x + \beta y + x^2 D_x + y^2 D_y) B_{n,o}^{(\alpha,\beta)}(x, y),$$

Indeed, by (8.9),

$$\begin{aligned} & (\alpha x + \beta y + x^2 D_x + y^2 D_y) B_{n,o}^{(\alpha,\beta)}(x, y) \\ &= (\alpha x + \beta y) \sum_{k=0}^n \binom{n}{k} (\alpha)_k (\beta)_{n-k} x^k y^{n-k} \\ &+ \sum_{k=0}^n k \binom{n}{k} (\alpha)_k (\beta)_{n-k} x^{k+1} y^{n-k} + \sum_{k=0}^n (n-k) \binom{n}{k} (\alpha)_k (\beta)_{n-k} x^k y^{n-k+1}. \end{aligned}$$

The coefficient of $x^k y^{n-k+1}$ on the right is equal to

$$\begin{aligned} & \alpha \binom{n}{k-1} (\alpha)_{k-1} (\beta)_{n-k+1} + \beta \binom{n}{k} (\alpha) (\beta)_{n-k} \\ & + (k-1) \binom{n}{k-1} (\alpha)_{k-1} (\beta)_{n-k+1} + (n-k) \binom{n}{k} (\alpha)_k (\beta)_{n-k} \\ & = \binom{n}{k-1} (\alpha)_k (\beta)_{n-k+1} + \binom{n}{k} (\alpha)_k (\beta)_{n-k+1} = \binom{n+1}{k} (\alpha)_k (\beta)_{n-k+1}. \end{aligned}$$

It follows from (8.11) that

$$(8.12) \quad B_{n,o}^{(\alpha,\beta)}(x,y) = (\alpha x + \beta y + x^2 D_x + y^2 D_y)^n \cdot 1.$$

9. When $\alpha = \beta = 0$, (8.1) reduces to

$$(9.1) \quad (xy)^n (D_x + D_y)^n = \sum_{k=0}^n B_{n,k}^{(0,0)}(x,y) (xy (D_x + D_y))^k,$$

while (8.3) becomes

$$(9.2) \quad B_{n=1,k}^{(0,0)}(x,y) = n(x+y) B_{n,k}^{(0,0)}(x,y) - n(n-1)xy B_{n-1,k}^{(0,0)}(x,y) \\ + B_{n,k-1}^{(0,0)}(x,y).$$

It is evident from (9.1) that

$$(9.3) \quad B_{n,o}^{(0,0)}(x,y) = 0 \quad (n > 0).$$

For brevity, put

$$b_{n,k} = \frac{1}{(n-1)!} B_{n,k}^{(0,0)}(x,y) \quad (n \geq 1).$$

Then (9.2) becomes

$$(9.4) \quad b_{n+1,k} - (x+y)b_{n,k} + xy b_{n-1,k} = \frac{1}{n} b_{n,k-1} \quad (k \geq 1).$$

For $k = 1$, (9.4) reduces to

$$(9.5) \quad b_{n+1,1} - (x+y)b_{n,1} + xy b_{n-1,1} = 0 \quad (n > 1).$$

The recurrence (9.5) implies

$$b_{n,1} = c_1 x^n + c_2 y^n,$$

where c_1, c_2 are constant. Since

$$b_{1,1} = 1, \quad b_{2,1} = x + y,$$

we get

$$(9.6) \quad b_{n,1} = \frac{x^n - y^n}{x - y} \equiv \sigma_n.$$

Next, for $k = 2$, we have

$$(9.7) \quad b_{n+1,2} - (x + y)b_{n,2} + xyb_{n-1,2} = \frac{n}{1} \sigma_n.$$

Since

$$b_{1,2} = b_{2,0} = 1,$$

(9.7) holds for $n \geq 1$. It follows that

$$(1 - (x + y)z + xyz^2) \sum_1^{\infty} b_{n+1,2} z^n = \sum_1^{\infty} \frac{n}{1} \sigma_n z^n.$$

Since

$$\frac{1}{1 - (x + y)z + xyz^2} = \sum_0^{\infty} \sigma_{n+1} z^n,$$

we get

$$b_{n+1,2} = \sum_{j=1}^n \frac{1}{j} \sigma_j \sigma_{n-j+1}.$$

Generally (9.4) implies

$$\sum_{n=k-1}^{\infty} (b_{n+1,k} - (x + y)b_{n,k} + xyb_{n-1,k}) z^n = \sum_{n=k-1}^{\infty} \frac{n}{1} b_{n,k-1} z^n,$$

that is,

$$(9.8) \quad \sum_{n=k-1}^{\infty} b_{n+1,k} z^n = \frac{1}{1 - (x + y)z + xyz^2} \sum_{n=k-1}^{\infty} \frac{n}{1} b_{n,k-1} z^n.$$

Therefore, as above, we get

$$(9.9) \quad b_{n+1,k} = \sum_{j=k-1}^n \frac{1}{j} b_{j,k-1} \sigma_{n-j+1}.$$

We may rewrite (9.9) in the form

$$(9.10) \quad b_{n+k,k} = \sum_{j=0}^n \frac{1}{j+k-1} b_{j+k-1,k-1} \sigma_{n-j+1}.$$

Using this formula we get

$$b_{n+3,3} = \sum_{0 \leq i \leq j \leq n} \frac{1}{(i+1)(j+2)} \sigma_{i+1} \sigma_{j-i+1} \sigma_{n-j+1},$$

$$b_{n+4,4} = \sum_{0 \leq i \leq j \leq k \leq n} \frac{1}{(i+1)(j+2)(k+3)} \sigma_{i+1} \sigma_{j-i-1} \sigma_{k-i+1} \sigma_{n-k+1}.$$

and so on

We may also mention an operational formula for $b_{n,k}$. Define the operator D_x^{-1} by means of

$$D_x^{-1} f(z) = \int_0^z f(t) dt.$$

Thus

$$\sum_{n=1}^{\infty} \frac{1}{n} \sigma_n z^n = D_x^{-1} \sum_{n=0}^{\infty} \sigma_{n+1} z^n = D_x^{-1} \frac{1}{(1-xz)(1-zy)}.$$

By (9.7)

$$\sum_{n=1}^{\infty} b_{n+1,2} z^n = \frac{1}{(1-xz)(1-yz)} D_x^{-1} \frac{1}{(1-xz)(1-yz)},$$

so that

$$\sum_{n=1}^{\infty} \frac{1}{n+1} b_{n+1,2} z^n = [D_x^{-1} (1-xz)^{-1} (1-yz)^{-1}]^2 \cdot 1.$$

At the next stage we get

$$\sum_{n=2}^{\infty} b_{n+1,3} z^n = [(1-xz)^{-1} (1-yz)^{-1} D_x^{-1}]^2 (1-xz)^{-1} (1-yz)^{-1}$$

The general formula is

$$(9.11) \quad \sum_{n=k-1}^{\infty} b_{n+1,k} z^n = [(1-xz)^{-1} (1-yz)^{-1} D_x^{-1}]^{k-1} (1-xz)^{-1} (1-yz)^{-1} \quad (k \geq 1).$$

10. A generating function for $B_{n,k}^{(\alpha,\beta)}(x,y)$ in the general case can be found in the following way. It follows from (8.5) that

$$(10.1) \quad \sum_{n=0}^{\infty} \frac{z^n}{n!} \sum_{k=j}^n (-1)^{k-j} C_{n,k}^{(\alpha,\beta)}(x,y) B_{k,j}^{(\alpha,\beta)}(x,y) = \frac{z^j}{j!}.$$

By (7.4) we have

$$\sum_{n=k}^{\infty} \frac{1}{n!} C_{n,k}^{(\alpha,\beta)}(x,y) = \frac{1}{k!} F^k(x,y) \Phi_{\alpha,\beta}(x,y).$$

Since $C_{n,k}^{(\alpha,\beta)}(x,y)$ is homogeneous of degree $n-k$, this implies

$$\sum_{n=k}^{\infty} \frac{z^n}{n!} C_{n,k}^{(\alpha,\beta)}(x,y) = \frac{z^k}{k!} F^k(xz,yz) \Phi_{\alpha,\beta}(xz,yz).$$

Thus (10.1) becomes

$$\begin{aligned} \frac{z^j}{j!} &= \sum_{k=j}^{\infty} (-1)^{k-j} B_{k,j}^{(\alpha,\beta)}(x,y) \sum_{n=k}^{\infty} \frac{z^n}{n!} C_{n,k}^{(\alpha,\beta)}(x,y) \\ &= \Phi_{\alpha,\beta}(xz,yz) \sum_{k=j}^{\infty} (-1)^{k-j} B_{k,j}^{(\alpha,\beta)}(x,y) \frac{z^k F^k(xz,yz)}{k!}. \end{aligned}$$

Multiplying by v^j and summing over j , we get

$$(10.2) \quad e^{xv} = \Phi_{\alpha,\beta}(xz,yz) \sum_{k=0}^{\infty} \sum_{j=0}^k (-1)^{k-j} B_{k,j}^{(\alpha,\beta)}(x,y) \frac{z^k F^k(xz,yz)}{k!} v^j.$$

Consider the equation

$$(10.3) \quad z F(xz,yz) = u,$$

that is,

$$\frac{e^{xz} - e^{yz}}{x e^{yz} - y e^{xz}} = u.$$

This reduces to

$$e^{(x-y)z} = \frac{1-xu}{1-yu},$$

so that

$$(10.4) \quad z = \frac{1}{x-y} \log \frac{1+xu}{1-yu} = \sum_1^{\infty} (-1)^{n-1} \frac{1}{n} \sigma_n u^n,$$

where as above

$$\sigma_n = (x^n - y^n)/(x - y).$$

Since

$$\begin{aligned} \Phi_{\alpha, \beta}(xz, yz) &= (1 + xzF(xz, yz))^\alpha (1 + yzF(xz, yz))^\beta \\ &= (1 + xu)^\alpha (1 + yu)^\beta, \end{aligned}$$

(10.2) becomes

$$(10.5) \quad e^{zv} = (1 + xu)^\alpha (1 + yu)^\beta \sum_{k=0}^{\infty} \frac{u^k}{k!} \sum_{j=0}^k (-1)^{k-j} B_{k,j}^{(\alpha, \beta)}(x, y) v^j.$$

But, by (10.4),

$$e^{zv} = \left(\frac{1 + xu}{1 + yu} \right)^{v/(x-y)},$$

so that (10.5) may be replaced by

$$(10.6) \quad \sum_{k=0}^{\infty} \frac{u^k}{k!} \sum_{j=0}^k B_{k,j}^{(\alpha, \beta)}(x, y) (x - y)^j v^j = \left(\frac{1 - xu}{1 - yu} \right)^{-v} (1 - xu)^{-\alpha} (1 - yu).$$

In particular, for $v = 0$, (10.6) reduces to

$$(10.7) \quad \sum_{k=0}^{\infty} \frac{k!}{u^k} B_{k,0}^{(\alpha, \beta)}(x, y) = (1 - xu)^{-\alpha} (1 - yu)^{-\beta},$$

which is evidently in agreement with (8.10).

For $\alpha = \beta = 0$, (10.6) becomes

$$(10.8) \quad \sum_{k=0}^{\infty} \frac{u^k}{k!} \sum_{j=0}^k B_{k,j}^{(0,0)}(x, y) (x - y)^j v^j = \left(\frac{1 - xu}{1 - yu} \right)^{-v}.$$

Since

$$\left(\frac{1 - xu}{1 - yu} \right)^{-v} = \sum_{r=0}^{\infty} \frac{(v)_r}{r!} x^r u^r \sum_{s=0}^{\infty} \frac{(-v)_s}{s!} y^s v^s,$$

we get

$$(10.9) \quad \sum_{j=0}^k B_{k,j}^{(0,0)}(x, y) (x - y)^j v^j = \sum_{r=0}^k \binom{k}{r} (-v)_r (v)_{k-r}.$$

The general result is only slightly more complicated, namely

$$(10.10) \quad \sum_{j=0}^k B_{k,j}^{(\alpha, \beta)}(x, y) (x - y)^j v^j = \sum_{r=0}^k \binom{k}{r} (\alpha + v)_r (\beta - v)_{k-r}.$$

It follows from (10.6), (10.7) and (10.8) that

$$\sum_{j=0}^k B_{k,j}^{(\alpha,\beta)}(x,y) v^j = \sum_{r=0}^k \binom{k}{r} B_{k-r,0}^{(\alpha,\beta)}(x,y) \sum_{j=0}^r B_{r,j}^{(0,0)}(x,y) v^j$$

and therefore

$$(10.11) \quad B_{k,j}^{(\alpha,\beta)}(x,y) = \sum_{r=0}^k B_{r,j}^{(0,0)}(x,y) B_{k-r,0}^{(\alpha,\beta)}(x,y).$$

Comparing coefficients of v^j on both sides of (10.8), we get

$$\sum_{k=j}^{\infty} \frac{u^k}{k!} B_{k,j}^{(0,0)}(x,y) = \frac{(-1)^j}{j!} \left(\log \frac{1-xu}{1-yu} \right)^j.$$

Differentiation with respect to u gives

$$\begin{aligned} & \sum_{k=j-1}^{\infty} \frac{u^k}{k!} B_{k+1,j}^{(0,0)}(x,y) \\ &= \frac{(-1)^{j-1}}{(j-1)!} (x-y)^{-j+1} \left(\log \frac{1-xu}{1-yu} \right)^{j-1} \frac{1}{(1-xu)(1-yu)} \end{aligned}$$

which is equivalent to (9.8).

REFERENCES

1. L. CARLITZ, *Eulerian numbers and polynomials*, *Mathematics Magazine*, vol. 33 (1959), pp. 247-260.
2. L. CARLITZ, *Enumeration of sequences by rises and falls: a refinement of the Simon Newcomb problem*, *Duke Mathematical Journal*, vol. 39 (1972), pp. 267-280.
3. L. CARLITZ and RICHARD SCOVILLE, *Generalized Eulerian numbers: combinatorial applications*, *Journal für die reine und angewandte Mathematik*, to appear.
4. J. F. DILLON and D. P. ROSELLE, *Simon Newcomb's problem*, *SIAM Journal on Applied Mathematics*, vol. 17 (1969), pp. 1086-1093.
5. D. FOATA and M. P. SCHUTZENBERGER, *Theorie Geometrique des Polynomes Euleriens*, *Lecture Notes in Mathematics*, 138, Springer-Verlag, Berlin, Heidelberg, New York, 1970.
6. JOHN RIORDAN, *An Introduction to Combinatorial Analysis*, Wiley, New York, 1958.
7. D. P. ROSELLE, *Permutations by number of rises and successions*, *Proceedings of the American Mathematical Society*, vol. 19 (1968), pp. 8-16.