

## Report on Twisted Sums of Banach Spaces

FÉLIX CABELLO AND JESÚS M.F. CASTILLO

*Dpto. de Matemáticas, Univ. de Extremadura, 06071-Badajoz, Spain*  
*e-mail: castillo@ba.unex.es*

AMS Subject Class. (1991): 46B20

Received October 14, 1996

### 1. INTRODUCTION

This note is to report some of the advances obtained as a follow-up of the book [2] on the topic of twisted sums of Banach spaces. Since this announcement is not longer enough to contain the theory being developed, we submit the interested reader to [2] and to [1], where full details and proofs shall appear.

### 2. BASICS ON TWISTED SUMS

A twisted sum of Banach spaces  $Y$ ,  $Z$  is a short exact sequence

$$0 \rightarrow Y \rightarrow X \rightarrow Z \rightarrow 0.$$

The open mapping theorem implies that  $Y$  is then a closed subspace of  $X$  and  $X/Y = Z$ .

The classical theory of Kalton-Peck [5] describes twisted sums in terms of homogeneous maps  $F : Z \rightarrow Y$  satisfying

$$\|F(x + y) - F(x) - F(y)\| \leq K (\|x\| + \|y\|),$$

which are called quasi-linear maps.

Given a twisted sum, the quasi-linear map that defines it can be obtained as the difference  $B - L$  between a bounded homogeneous and a linear selection for the quotient map. Conversely, if  $F : Z \rightarrow Y$  is a quasi-linear map, the product space  $Y \times Z$  endowed with the quasi-norm

$$\|(y, z)\| = \|y - F(z)\| + \|z\|$$

is denoted  $Y \oplus_F Z$  and provides a twisted sum

$$0 \rightarrow Y \rightarrow Y \oplus_F Z \rightarrow Z \rightarrow 0.$$

For instance, the direct sum  $Y \oplus Z$  is the twisted sum space provided by any linear map  $Z \rightarrow Y$ . When a twisted sum is equivalent to the direct sum (i.e. when  $Y$  is complemented in  $X$ ) we also say that the twisted sum splits. Kalton and Peck [5] showed that two twisted sums  $Y \oplus_F Z$  and  $Y \oplus_G Z$  are equivalent if and only if for some linear map  $L : Z \rightarrow Y$

$$\text{dist}(F - G, L) < +\infty.$$

In particular, the twisted sum defined by  $F$  splits if and only if  $\text{dist}(F, L) < +\infty$  for some linear map  $L : Z \rightarrow Y$ .

Observe that the twisted sum space  $Y \oplus_F Z$  is not necessarily locally convex: in fact, the expression  $\|(y, z)\| = \|y - F(z)\| + \|z\|$  is just a quasi-norm. Kalton [4] proved that when  $Y$  and  $Z$  are  $B$ -convex Banach spaces then  $Y \oplus_F Z$  is a Banach space too.

### 3. NEWS ON TWISTED SUMS

The problem of when  $Y \oplus_F Z$  is a Banach space can be completely solved (see [2]). Let us call a homogeneous map  $F : Z \rightarrow Y$  0-linear if whenever  $\sum_{i=1}^n x_i = 0$  then

$$\left\| \sum_{i=1}^n F(x_i) \right\| \leq K \sum_{i=1}^n \|x_i\|$$

for some constant  $K > 0$  independent of the points  $x_i$ .

**THEOREM.** *The expression  $\|(y, z)\| = \|y - F(z)\| + \|z\|$  is equivalent to a norm if and only if  $F$  is 0-linear.*

(To this, one should add the following result of D. Yost: the expression  $\|(y, z)\| = \|y - F(z)\| + \|z\|$  is itself a norm if and only if  $F$  is pseudo-linear in the sense that  $\|F(x + y) - F(x) - F(y)\| \leq \|x\| + \|y\| - \|x + y\|$ ).

### 4. THE NONLINEAR HAHN-BANACH THEOREM

Therefore, by the Hahn-Banach theorem, if  $F : Z \rightarrow \mathbb{R}$  is a 0-linear map, the twisted sum it defines splits and then, by the criterium of Kalton and Peck, for some linear map  $L : Z \rightarrow \mathbb{R}$

$$\text{dist}(F, L) < +\infty.$$

THEOREM. *An explicit construction of a linear map at finite distance of a 0-linear given map  $F : Z \rightarrow \mathbb{R}$ .*

#### 5. SOBCZYK'S THEOREM KALTON'S WAY

Sobczyk's theorem asserts that  $c_0$  is complemented in any separable Banach space containing it. In our language, this means that every 0-linear map  $F : Z \rightarrow c_0$  with  $Z$  separable admits a linear map  $L : Z \rightarrow c_0$  at finite distance.

THEOREM. *An explicit construction of a linear map  $L : Z \rightarrow c_0(I)$  at finite distance of a 0-linear given map  $F : Z \rightarrow c_0(I)$ , when  $Z$  is separable.*

#### 6. NONLINEAR DUALITY

It is well-known that if  $0 \rightarrow Y \rightarrow X \rightarrow Z \rightarrow 0$  is an exact sequence then  $0 \rightarrow Z^* \rightarrow X^* \rightarrow Y^* \rightarrow 0$  is also exact. This means that if  $F : Z \rightarrow Y$  is a 0-linear map there should be

THEOREM. *An explicit method to construct the adjoint 0-linear map  $F^* : Y^* \rightarrow Z^*$ .*

Knowing the form of  $F^*$  one can prove that the space  $Z_2$  of Kalton-Peck [5] is isomorphic to its dual.

Derived from this we consider two related

#### 7. THREE-SPACE PROBLEMS ON DUALITY

It is an open problem to know if "being a dual space" is a three-space property, in the sense that given an exact sequence  $0 \rightarrow Y^* \rightarrow X \rightarrow Z^* \rightarrow 0$  the space  $X$  must be a dual space. We construct

THEOREM. *An exact sequence  $0 \rightarrow Y^* \rightarrow X \rightarrow Z^* \rightarrow 0$  that is not a dual sequence,*

which solves a question that goes back to Vogt [7]. Moreover,

THEOREM. *If  $0 \rightarrow Y^* \rightarrow X \rightarrow R \rightarrow 0$  is an exact sequence, where  $R$  is reflexive, then it is a dual sequence.*

Related to this is the question: is the property of being complemented in its bidual a three-space property? This question is also open. A simplification of the preceding argument shows:

**THEOREM.** *If  $0 \rightarrow Y \rightarrow X \rightarrow R \rightarrow 0$  is an exact sequence where  $Y$  is complemented in its bidual and  $R$  is reflexive then  $X$  is complemented in its bidual (see also [3]).*

In some cases, there is a positive answer:

**THEOREM.** *If  $0 \rightarrow Y \rightarrow X \rightarrow Z \rightarrow 0$  is an exact sequence where  $Y$  is complemented in some dual space and  $Z$  is an  $\mathcal{L}_1$ -space then the sequence splits.*

This contains an old result of Lindenstrauss [6].

#### REFERENCES

- [1] CABELLO, F. AND CASTILLO, J.M.F., Twisted sums of Banach spaces, preprint.
- [2] CASTILLO, J.M.F. AND GONZÁLEZ, M., "Three-space problems in Banach space theory," monograph submitted.
- [3] DÍAZ, J.C., DIEROLF, S., DOMANSKI, P. AND FERNÁNDEZ, C., On the three-space problem for dual Fréchet spaces, *Bull. Acad. Polon. Sci.*, **40** (1992), 221–224.
- [4] KALTON, N., The three-space problem for locally bounded  $F$ -spaces, *Comp. Math.*, **37** (1978), 243–276.
- [5] KALTON, N. AND PECK, N.T., Twisted sums of sequence spaces and the three-space problem, *Trans. Amer. Math. Soc.*, **255** (1979) 1–30.
- [6] LINDENSTRAUSS, J., On a certain subspace of  $l_1$ , *Bull. Acad. Polon. Sci.*, **12** (1964), 539–542.
- [7] VOGT, D., Lectures on projective spectra of  $(DF)$ -spaces, preprint.

