

EMBEDDING TORSIONLESS MODULES IN PROJECTIVES

CARL FAITH

Abstract

In this paper we study a condition right FGTF on a ring R , namely when all finitely generated torsionless right R -modules embed in a free module. We show that for a von Neuman regular (VNR) ring R the condition is equivalent to every matrix ring R_n is a Baer ring; and this is right-left symmetric. Furthermore, for any Utumi VNR, this can be strengthened: R is FGTF iff R is self-injective.

Introduction

When all injective modules over a ring R embed in a free R -module, the ring R must be quasi-frobenius, ($=$ QF) and conversely. In this case every right or left R -module embeds in a projective R -module, and furthermore, the one-sided condition implies the two-sided condition ([F-W]). The two-sided condition that all cyclic R -modules embed in projectives implies R is QF, but here the one-sided condition is not sufficient.

Related to these rings are right IF rings initiated by Jain [J], Colby [Co], Damiano [D], and Würfel [W]. A ring R is **right IF** if every (injective) right module embeds in a flat module, i.e. if every injective right R -module is flat. These rings were characterized by Colby and Würfel in the one-sided case by the property that all finitely presented right R -modules embed in projective modules. Here, again, right does not imply the left condition ([Co]). However, two-sided IF are coherent (l.c.). A number of characterizations of IF rings are summarized in Section 4.

In this paper we study a condition that is considerably weaker, namely **right FGTF**: all finitely generated torsionless right R -modules embed in a projective equivalently in a free right R -module. We let **right FPTF** denote the same condition for finitely presented torsionless right R -modules.

We now introduce a concept that is germane to FGTF: A ring R is a **right \star -ring** (star ring) provided that every finitely generated right R -module has finitely generated dual.

Easy exercises establish that any left Noetherian ring is right \star , and that any right \star -ring is right FGTF. We establish here that over any von Neumann

regular (= VNR) ring R that (right or left) FGTF and \star rings are equivalent concepts. This relates to a theorem of Kobayashi that states that any right self-injective VNR is a right \star -ring. Furthermore, we establish the converse to Kobayashi's theorem for Utumi VNR's. This generalizes his own converse theorem for commutative VNR's to VNR's with (2-sided) self-injective maximal right quotient ring $Q_{max}^r(R)$, e.g. to Abelian VNR's. As a corollary we show that over an Abelian VNR ring R , that every matrix ring R_n is Baer iff R is self-injective. The proof requires two theorems of Utumi: (1) on the left self-injectivity of the maximal right quotient ring $Q_{max}^r(R)$ of a non-singular ring R ; (2) on self-injectivity of continuous $n \times n$ matrix rings, $n \geq 2$.

We also show by example that the converse to Kobayashi's theorem fails for all non-Utumi VNR's.

In Section 5, we relate IF, FGTF and \star rings.

1. Preliminaries

For a subset X of R , we let X^\perp denote the right, ${}^\perp X$ the left, annihilator in R . Thus, a right ideal I is an annihilator iff $I = X^\perp$ for $X \subseteq R$. **Annulet** is a variant term for annihilator. Then, I is said to be **finitely annihilated** (= **FA**) provided that $I = X^\perp$ for a finite set X . If a ring satisfies the acc on annihilator left ideals (= R is a \perp acc ring), then every right annihilator of R is FA by a theorem of [F1].

A right R -module is **torsionless** if M has the equivalent properties: (T1) M embeds in a direct product of copies of R ; (T2) M canonically embeds in its bidual module M^{**} , (T3) For each nonzero $x \in M$ there exists $f \in M^* = \text{Hom}_R(M, R)$ so that $f(x) \neq 0$.

Moreover, for any ring A , a cyclic right module A/I is:

- (1) torsionless $\Leftrightarrow I = X^\perp$ for a subset X of A (= I is a right annihilator)
- (2) embeddable in a free module $\Leftrightarrow I = X^\perp$ for $|X| < \infty$ (= I is FA).

We also record the fact mentioned in the abstract:

1.0. Proposition. *A right \star -ring is right FGTF.*

Proof: Trivial. (See, e.g. [F2].) ■

1.1. Theorem. *A ring R is right FGTF iff each annihilator right ideal of the $n \times n$ matrix ring R_n is FA for every n .*

Proof: The proof of this Folkloric theorem is by "Morita theory", i.e. using the fact that under the Morita equivalence

$$\Theta \left\{ \begin{array}{l} \text{mod} - R \rightsquigarrow \text{mod} - R_n \\ X \longrightarrow X^n \end{array} \right.$$

that n -generated right R -modules map onto cyclic right R_n -modules. Furthermore, a submodule X of free R -module, say $X \subseteq R^n$ maps onto a principal right ideal of R_n under Θ . Similarly, Θ preserves torsionless modules. ■

1.2. Corollary. *If a ring R embeds in left Noetherian ring A , then R is right FGTF.*

Proof: Then R_n embeds in a left Noetherian ring A_n , for every n , and thus satisfies the acc on annihilator left ideals. By [F1], then each annihilator right ideal is FA, so R is right FGTF. ■

2. Semi-Continuous and Baer Rings

A ring R is said to be **right semi-continuous** ([F3]) if R has the property:

right SC: every right ideal is an essential submodule of one generated by an idempotent.

In [F3] we indicated that this is equivalent to:

right CS: every complement right ideal is generated by an idempotent.

A ring R is **right continuous** provided that R is right SC and every right ideal I which is isomorphic to a right ideal generated by an idempotent is itself a right ideal generated by an idempotent.

A ring R is **VNR** (= von Neumann regular) iff each finitely generated right ideal is generated by an idempotent. (This is right left symmetric.) Obviously a VNR is right continuous iff R is right SC.

A VNR ring R is **Abelian** or **strongly regular** iff every idempotent in R is central. Moreover, an idempotent $e \in R$ is **Abelian** if eRe is an Abelian VNR.

2.1. Theorem. *If R is a right continuous VNR ring, then R is right self-injective under any of the following assumptions:*

- (1) R has no nonzero Abelian idempotents.
- (2) $M_n(R)$ is right continuous for any $n > 1$ (Utumi [U₁]).
- (3) R is directly indecomposable qua ring (e.g., R prime).
- (4) Every primitive factor ring is Artinian.

See [G2] for proofs.

A ring R is a **Baer ring** if every right annihilator is generated by an idempotent. (This is left-right symmetric.) By the Johnson-Utumi theory of maximal quotient rings (see [F4, chap. 19.]) every annihilator right ideal of a right non-singular ring R is a complement right ideal and, thus, is a direct summand when R is right self-injective, a fact that we record:

Theorem of R.E. Johnson and Y. Utumi. *The maximal right quotient ring $Q_{\max}^r(R)$ of a right nonsingular ring R is a right self-injective VNR ring, hence a right continuous Baer ring.*

2.2. Theorem. *A VNR ring R is right FGTF iff every R_n is a Baer ring for every $n \geq 1$.*

Proof: This follows from Theorem 1.1, since every finitely generated one-sided ideal of a VNR ring is generated by an idempotent, and hence then so is its annihilator. ■

2.3. Theorem. *For a VNR ring, the f.a.e.:*

- (1) R is right FGTF.
- (2) R is left FGTF.
- (3) R is a right \star -ring.
- (4) R is a left \star -ring.
- (5) R_n is a Baer ring $\forall n$.

Proof: Since "Baer ring" is left-right symmetric, (1) \Leftrightarrow (2) \Leftrightarrow (5) by Theorem 2.2. Moreover, (3) \Rightarrow (1) by Theorem 1.0, and (1) \Rightarrow (4) by Theorem 2.2 and Camillo's Theorem [C] cited below. (See Remark 2.5.A.) The remaining implications follow by symmetry. ■

An R -module is **coherent** if every finitely-generated submodule is finitely presented.

2.4 Theorem ([C]). *For any ring R the f.a.e.:*

- (1) R is right Π -coherent, i.e. any product of copies of R is a coherent right R -module.
- (2) R is a left \star -ring.
- (3) Annihilator right ideals of $M_n(R)$ are finitely generated, $\forall n \geq 1$.

2.5A Remark. Any right coherent right FGTF ring R is right Π -coherent, hence a left \star -ring, whence left FGTF.

Thus by Camillo's theorem:

2.5B Corollary. *A left coherent right Π -coherent ring is left Π -coherent.*

Proof: R is left \star , hence left FGTF, so 2.5A applies. ■

2.5C Corollary. *A left coherent left \star -ring is left Π -coherent.*

2.5D Remark. We can easily deduce Theorem 1.0 from Theorem 2.4, using Theorem 1.1, since (3) of Theorem 2.4 implies the condition of Theorem 1.1.

Proof: This follows from Theorem 1.0, and the fact that any finitely generated submodule of a free module is coherent over a coherent ring. ■

3. Utumi Rings

As stated *supra* the Johnson-Utumi theorem in a right non-singular ring R , every right annulet is a complement right ideal. Utumi [U₂] considered the converse condition:

(right U) Every right complement is a right annulet.

We say R is **right Utumi** if R satisfies this condition, and R is **Utumi** if R is right & left Utumi. A **non-singular ring** R is a right and left non-singular ring.

Utumi's theorem. ([U₂]). *In a non-singular ring R , the f.a.e.:*

- (1) *The maximal right quotient ring $Q_{max}^r(R)$ is left self-injective.*
- (2) $Q_{max}^r(R) = Q_{max}^l(R)$
- (3) *R is Utumi.*

The next proposition is almost obvious.

3.1 Proposition. *A Baer VNR right Utumi ring R is right continuous.*

Proof: R is right SC (= CS), since every complement right ideal is generated by an idempotent. ■

3.2 Theorem. *For an Utumi-VNR the f.a.e.:*

- (1) *R is right FGTF.*
- (2) *R is left FGTF.*
- (3) *R is right self-injective.*
- (4) *R is left self-injective.*

Proof: It follows from Proposition 3.1 and Theorem 2.2 that $M_n(R)$ is right continuous $\forall n \geq 1$ when R is left or right FGTF, hence from Theorem 2.1 that $M_n(R)$ is (right and left) self-injective. Thus (1) \Rightarrow (3), and (2) \Rightarrow (3).

If R is right self-injective then $M_n(R)$ is right continuous, since $M_n(R)$ is right self-injective. Thus, $M_n(R)$ is a Baer ring $\forall n \geq 1$. Then R is right and left FGTF by Theorem 1.1, so (3) \Rightarrow (1). Then R is left self-injective by (2) \Rightarrow (3), hence therefore (3) \Rightarrow (4). Using Theorem 2.2, this completes the Proof. ■

3.3 Corollary. *An Abelian VNR ring R is right FGTF iff R is self-injective.*

3.4. Example. Any right but not left injective ring Q , e.g. the maximal right quotient ring $Q_{max}^r(R)$ of a VNR ring R that is not Utumi, is a left \star -ring that is not left self-injective.

It follows from Kobayashi's theorem that R is right \star -ring and from Theorem 2.3 that R is a left \star -ring. ■

4. If Rings

A number of these results are related to theorems of Gómez-Pardo and González, Goodearl, and Levy on when do finitely presented cyclic [GP-G], finitely generated nonsingular ([G1]), or finitely generated torsion-free modules respectively embed in free modules (see [F2]).

Moreover:

4.1. Theorem. (Colby [Co], Würfel[W])

All finitely presented right R -modules embed in a free module iff R is a right IF ring.

4.2. Theorem. ([J], [Co], [W], and [GP-G].)

The f.a.e.c.'s on a ring R .

(IF1) R is right and left IF.

(IF2) R is (right & left) coherent and every finitely generated one-sided ideal is an annihilator.

(IF3) Annihilation defines a duality between finitely generated one-sided ideals.

(IF4) R is coherent and every flat module (either side) is FP-injective.

(IF5) The classes of flat left and \aleph_0 -injective left R -modules are equal.

A ring R is **right (F)GF** if every (finitely generated) right R -module embeds in a free right R -module. Every quasi-Frobenius (QF) ring is both right and left GF.

In this connection the following theorem is of interest.

4.3. Theorem. (Jain [J], Rutter [R]). *Any right FGF ring is right IF.*

Also of interest here are certain other theorems of Jain [J] which are summarized in [F2, p.35, Theorem 4.2 and Corollaries].

In [F2] we discussed various conditions that implied that right FGF rings are quasi-Frobenius (QF), including the assumption that R is left and right FGF. The general FGF problem, however, remains open. (However, see [Me].)

If F is a right cogenerator ring, then every right R -module is torsionless, hence R is then FGF iff FGTF. Whether R is necessarily QF is an open question, so the classification FGTF problem will remain open until the FGF problem for right cogenerator rings is settled. (Note, however, by a theorem of Onodera, any right and left cogenerator ring is self-injective and right self-injective right FGF rings are QF (see [F2] and [F5]).

5. If \star -Rings

An element $r \in R$ is **regular** if r is neither a right nor left zero divisor of R . Let R^* denote the set of zero divisors of R . A right R -module M is

torsion-free if:

$$xr = 0 \Rightarrow x = 0 \forall x \in M, r \in R^*$$

A ring R is **right Ore** if and only if R has a classical right quotient ring

$$Q = Q_{Cl}^r(R) = \{a/b | a \in R, b \in R^*\}$$

where addition and multiplication in Q is canonical.

It is evident that over Q every right R -module is torsion-free.

Consider the property **right f.g.t.f.** = all finitely generated torsion-free right R -modules embed in a free module.

We now come to Levy's theorems.

Levy's first Theorem. *If R is a semi-prime right Ore ring then R is right f.g.t.f. iff R is a left Ore ring. In this case $Q_{Cl}^l(R) = Q_{Cl}^r(R)$ is semisimple.*

By a remark above, one sees that a ring $Q = Q_{Cl}^r(R)$ right FGF iff Q is right f.g.t.f. In this case, by [F-W] we have:

5.1. Theorem. ([F2]) *If R is left and right Ore, then R is right and left f.g.t.f. iff Q is QF.*

5.2. Corollary. ([F2]) *A commutative ring R is f.g.t.f. iff $Q_{Cl}(R)$ is QF.*

Since any torsionless module is torsion-free, one concludes that any commutative f.g.t.f. ring R , hence any R with $Q_{Cl}(R)$ QF, is right FGTF.

5.3. Remark. It follows from 5.2 that an integral domain R that is not coherent is an example of an FGTF ring that is not \star .

5.4. Theorem. *The f.a.e.c.'s on a ring R .*

- (1) R is a right and left FGTF and all finitely generated one-sided ideals are annulets.
- (2) R is a right and left IF \star -ring.
- (3) R is an IF and FGTF ring.

Proof: (2) \Rightarrow (1) by Theorem 1.0 and Theorem 4.2, (IF2).

(1) \Rightarrow (2). All annihilator in any matrix ring R_n right or left ideals are FA by Theorem 1.1. Now let I be any right annulet. Now $L = {}^\perp I$ is FA, say $L = {}^\perp I_1$, for I_1 a finitely generated right ideal, which by hypothesis is an annulet. Thus, $L^\perp = I = I_1$ is finitely generated. By (IF3) of Theorem 4.2, R is an IF ring. By (IF2), R is coherent, hence R is a \star -ring by Theorem 2.4 and Remark 2.5A.

Obviously (2) \Rightarrow (3). Moreover (3) \Rightarrow (2) since IF implies R is coherent so Remark 2.5A applies. ■

6. Problems

1. Characterize the condition that all annihilator right ideals of $M_n(R)$ are finitely generated (for all $n \geq 1$) ideal-theoretically in R .

2. Same question for $acc\perp$ ($dcc\perp$) in $M_n(R)$.

In [C], Camillo raises the question:

3. (Question of [C]): If R is a (2-sided) Noetherian ring, is the polynomial ring $R[X]$ Π -coherent for any set X ?

Camillo verifies 3. for semiprime R and, in fact, for any R embeddable in an Artinian ring. (A semiprime Noetherian ring is Goldie and has Artinian classical ring of quotients.) However, as proved by Dean and Stafford [D-S], not every Noetherian ring can be embedded in an Artinian ring. Nevertheless, using a theorem of [C-G], Camillo [C] verifies 3. for any Noetherian ring R that is an algebra over a non-denumerable field.

4. Characterize right FGTF rings, equivalently, by Theorem 1.1, characterize when every annihilator right ideal of R_n , $n \geq 1$, is FA. (This characterization should be ideal - theoretically in R .)

References

- [B] H. BASS, Finitistic dimension and a homological generalization of semi-primary rings, *Trans. Amer. Math. Soc.* **95** (1960), 466-88.
- [Be] S.K. BERBERIAN, "Baer *-rings," Springer Verlag, Berlin-Heidelberg-New York, 1972.
- [C] V. CAMILLO, On the coherence of polynomial rings, *J. Algebra* (to appear).
- [C-G] V. CAMILLO AND R. GURALNIK, Polynomial rings over Goldie rings are often Goldie, *Proc. Amer. Math. Soc.* **98** (1986).
- [Co] R.R. COLBY, Rings which have flat injectives, *J. Alg.* **35** (1975), 239-252.
- [D] R.F. DAMIANO, Coflat rings and modules, *Pac. J. Math.* **81** (1979), 349-369.
- [D-S] C. DEAN AND J.T. STAFFORD, A non-embeddable Noetherian ring, *J. Alg.* **115** (1988), 175-181.
- [F1] C. FAITH, Rings with ascending condition on annihilators, *Nagoya Math. J.* **27** (1966), 179-191.
- [F2] C. FAITH, "Embedding modules in projectives, in Advances in Non-commutative Ring Theory," LNM **951**, 1982, pp. 21-39.
- [F3] C. FAITH, The maximal regular ideal of self-injective and continuous rings splits off, *Arch. Math.* **44** (1985), 511-525.

- [F4] C. FAITH, "Algebra II: Ring Theory," Springer-Verlag, Berlin-Heidelberg-New York, 1976.
- [F5] C. FAITH, When self-injective rings are QF: a report on a problem (preprint).
- [F-W] C. FAITH AND E.A. WALKER, Direct sum representations of injective modules, *J. Alg.* **5** (1967), 203-221.
- [GP-G] J.L. GÓMEZ-PARDO, AND N.R. GONZÁLEZ, On some properties of IF rings, *Quaestiones Math.* **5** (1983), 395-405.
- [G1] K.R. GOODEARL, Embedding non-singular modules in free modules, *J. Pure Appl. Algebra* **1** (1971), 275-279.
- [G2] K.R. GOODEARL, "Von Neuman Regular Rings," Pitman, London, San Francisco, and Melbourne, 1976.
- [K] S. KOBAYASHI, A note on regular self injective rings, *Osaka J. Math.* **21** (1984).
- [J] S. JAIN, Flat and FP-injectivity, *Proc. Amer. Math. Soc.* **41** (1973), 437-442.
- [Me] P. MENAL, On the endomorphism ring of a free module, *Pub. Mats. U. Autònoma de Barcelona (U.A.B.)* **27** (1982), 141-154.
- [S] B. STENSTROM, "Rings of Quotients," Springer-Verlag, Berlin, Heidelberg, New York, 1975.
- [U₁] Y. UTUMI, On continuous regular rings and semisimple self-injective rings, *Can. Math. J.* **12** (1960), 597-605.
- [U₂] Y. UTUMI, Of rings of which any one-sided quotient rings are two-sided, *Proc. Amer. Math. Soc.* **14** (1963), 141-147.
- [W] T. WÜRFEL, Über absolut reine Ringe, *J. Reine Angew. Math.* **262/263** (1973), 381-391.

The State University of New Jersey RUTGERS
Department of Mathematics at New Brunswick
Hill Center for the Mathematical Sciences
Busch Campus New Brunswick
New Jersey 08903
U.S.A.

Rebut el 17 de Gener de 1990