

THE p -PERIOD OF AN INFINITE GROUP

YINING XIA

Abstract

For Γ a group of finite virtual cohomological dimension and a prime p , the p -period of Γ is defined to be the least positive integer d such that Farrell cohomology groups $\hat{H}^i(\Gamma; M)$ and $\hat{H}^{i+d}(\Gamma; M)$ have naturally isomorphic p -primary components for all integers i and $Z\Gamma$ -modules M .

We generalize a result of Swan on the p -period of a finite p -periodic group to a p -periodic infinite group, i.e., we prove that the p -period of a p -periodic group Γ of finite vcd is $2LCM(|N(\langle x \rangle)/C(\langle x \rangle)|)$ if the Γ has a finite quotient whose a p -Sylow subgroup is elementary abelian or cyclic, and the kernel is torsion free, where $N(-)$ and $C(-)$ denote normalizer and centralizer, $\langle x \rangle$ ranges over all conjugacy classes of Z/p subgroups. We apply this result to the computation of the p -period of a p -periodic mapping class group. Also, we give an example to illustrate this formula is false without our assumption.

For Γ a group of virtual finite cohomological dimension (vcd) and a prime p , the p -period of Γ is defined to be the least positive integer d such that the Farrell cohomology groups $\hat{H}^i(\Gamma; M)$ and $\hat{H}^{i+d}(\Gamma; M)$ have naturally isomorphic p -primary components for all $i \in \mathbb{Z}$ and $Z\Gamma$ -modules M [3].

The following classical result for a finite group G was showed by Swan in 1960 [9].

Theorem (Swan).

- a) If a 2-Sylow subgroup of G is cyclic ($\neq \{1\}$), the 2-period of G is 2. If a 2-Sylow subgroup of G is a (generalized) quaternion group, the 2-period of G is 4.
- b) Suppose p an odd prime and a p -Sylow subgroup of the finite group G is cyclic ($\neq \{1\}$). Let S_p denote the p -Sylow subgroup and A_p the group of automorphisms of S_p induced by inner automorphism of G . Then the p -period of G is twice the order of A_p .

Remark.

The group A_p above is isomorphic to $N(S_p)/C(S_p)$, where $N(-)$ and $C(-)$ denote the normalizer and centralizer of S_p in G .

It is very natural to ask a question: If Γ is a p -periodic group of finite vcd , is a similar result still true? In other words, is it possible to describe the p -period of a p -periodic group Γ of finite vcd by an algebraic *non-homological* invariant of the group Γ itself?

In this paper, we generalize the result of Swan for a finite group to a p -periodic group Γ of finite vcd which has a finite quotient whose a p -Sylow subgroup is elementary abelian or cyclic, and the kernel is torsion-free, i.e., we prove that the p -period of a p -periodic group Γ of finite vcd is twice the least common multiple of $\{|N(\langle x \rangle)/C(\langle x \rangle)|\}$ in these two cases, where $\langle x \rangle$ ranges over all conjugacy classes of Z/p subgroups of Γ . On the other hand, we give a group Γ_0 of finite vcd whose only finite subgroup is a $Z/2$, but the 2-period of Γ_0 is greater than $2|N(Z/2)/C(Z/2)|$. Finally, an application will be made for calculating the p -period of a mapping class group.

The following four theorems are our main results of this paper.

Theorem 1. *Assume that Γ is p -periodic. If Γ has a normal subgroup of finite cohomological dimension so that the associated quotient is a finite group whose a p -Sylow subgroup is elementary abelian, then the p -period of Γ is twice the least common multiple of $\{|N(\langle x \rangle)/C(\langle x \rangle)|\}$, where $\langle x \rangle$ ranges over all conjugacy classes of Z/p subgroups of Γ .*

Theorem 2. *Let Γ be a group which has a normal subgroup of finite cohomological dimension so that the associated quotient is a finite group whose a p -Sylow subgroup is cyclic, then the p -period of Γ is twice the least common multiple of $\{|N(\langle x \rangle)/C(\langle x \rangle)|\}$, where $\langle x \rangle$ ranges over all conjugacy classes of Z/p subgroups of Γ .*

Theorem 3. *There is a group Γ_0 of finite vcd whose only finite subgroup is a $Z/2$, but the 2-period is greater than $2|N(Z/2)/C(Z/2)|$.*

Theorem 4. *If the mapping class group Γ_g is a p -periodic group and $g < p(p-1)/2$, then the p -period of Γ_g is $2\text{LCM}\{\text{gcd}(p-1, b_i)\}$, where $b_i \in B_{g,p}$ (cf. section 3).*

The rest of this paper is organized as follows. In section 1, we prove Theorems 1 and 2. In section 2, we provide an example illustrating Theorem 3. Finally in section 3, we give a formula for the calculation of the p -period of a p -periodic mapping class group Γ_g .

1. Proof of Theorems 1 and 2

Lemma 1.1. *Let $H = \langle x, y/x^p = 1, yxy^{-1} = x^r \rangle$, where $q = 0$ or $q \neq 0 \pmod{p}$. If d is the minimal positive integer such that $r^d = 1 \pmod{p}$, then the p -period of H equals $2d$.*

Proof: If $q \neq 0$, H is a finite group, the proof is immediate by Swan Theorem. Otherwise, if $q = 0$, H is infinite and we look at the short exact sequence $1 \rightarrow Z/p \rightarrow H \rightarrow Z \rightarrow 1$. The spectral sequence of Farrell cohomology associated to the exact sequence converges in the following way: $E_2^{i,j} = H^i(Z; \hat{H}^j(Z/p; Z)) \rightarrow \hat{H}^{i+j}(H; Z)$ [2]. This spectral sequence collapses since $H^i(Z; \hat{H}^j(Z/p; Z)) = 0$ when $i < 0$ or $i > 1$. Therefore, $1 \rightarrow \hat{H}^{n-1}(Z/p; Z)_Z \rightarrow \hat{H}^n(H; Z) \rightarrow \hat{H}^n(Z/p; Z)^Z \rightarrow 1$ is an exact sequence. By looking at the Z action on the subgroup $Z/p, u^d \in \hat{H}^{2d}(Z/p; Z)$ is an invariant element of the Z action on $\hat{H}^{2d}(Z/p; Z)$. Here u is a generator of $\hat{H}^2(Z/p, Z)$. Therefore, there exists an element $h \in \hat{H}^{2d}(H; Z)$ such that $\text{Res}(h) = u^d \neq 0$ on $\hat{H}^{2d}(Z/p; Z)$. By Brown-Venkov theorem [2] and $\hat{H}^{2kd}(H; Z) = Z/p, \hat{H}^{2kd+1}(H; Z) = Z/p, \hat{H}^i(H; Z) = 0$ for other i 's, the p -period of H is $2d$. ■

Lemma 1.2. *Let Z/p be a normal subgroup of a group Γ of finite vcd, and let M be a finite quotient of Γ with torsion free kernel. Then $\Gamma/C_\Gamma(Z/p) = N_\Gamma(Z/p)/C_\Gamma(Z/p) = N_M(Z/p)/C_M(Z/p) = M/C_M(Z/p)$. Here we still use Z/p to stand for the image of Z/p in M .*

Proof: Let $pr : \Gamma \rightarrow M$ be the natural projection map. The map pr maps $N_\Gamma(Z/p)$ onto $N_M(Z/p)$ and $C_\Gamma(Z/p)$ to $C_M(Z/p)$, so induced map $pr_* : N_\Gamma(Z/p)/C_\Gamma(Z/p) \rightarrow N_M(Z/p)/C_M(Z/p)$ is a well-defined surjective homomorphism. Let $\langle x \rangle = Z/p$, if $yxy^{-1} = x^r$, then $pr(y)xpr(y)^{-1} = x^r$, i.e., pr_* is an injective. ■

Lemma 1.3. *Suppose a group M contains a cyclic subgroup $Z/p^n \supset Z/p$ and $|N(Z/p^n)/C(Z/p^n)|$ is prime to p , then the homomorphism induced by inclusion $i_* : N(Z/p^n)/C(Z/p^n) \rightarrow N(Z/p)/C(Z/p)$ is injective.*

Proof: Notice $N(Z/p) \supset N(Z/p^n)$ and the inclusion i maps $C(Z/p^n)$ to $C(Z/p)$, i.e., the induced map by inclusion $i_* : N(Z/p^n)/C(Z/p^n) \rightarrow N(Z/p)/C(Z/p)$ is a well-defined homomorphism. Now let $\langle x \rangle = Z/p^n$, then $\langle x^{p^{n-1}} \rangle = Z/p$, if $y \in C(Z/p)$, $yxy^{-1} = x^k$, then $yx^{p^{n-1}}y^{-1} = x^{kp^{n-1}} = x^{p^{n-1}}$, so $(k-1)p^{n-1} = 0 \pmod{p^n}$, i.e., $k = 1 \pmod{p}$. Let $k = Ap^m + 1$, A is prime to p and $1 \leq m < n$, $k^d = 1 \pmod{p^n}$, d divides $p-1$

by assumption. Hence $k^d = (Ap^m + 1)^d = B + Adp^m + 1 = 1 \pmod{p^n}$, where p^{2m} divides B . This implies $Ad = 0 \pmod{p}$, a contradiction unless $A = 0$. ■

Lemma 1.4 (Swan) [9]. *Suppose the p -Sylow subgroup S_p of a finite group M is abelian. Let A_p be the group of automorphisms of S_p induced by inner automorphisms of M . Then an element $a \in H^i(S_p; Z)$ is stable if and only if it is fixed under the action of A_p on $H^i(S_p; Z)$.*

Proof: See [9]. ■

Proof of Theorem 1: A theorem of Brown [3, p. 293] states that if Γ is p -periodic, then $\hat{H}^*(\Gamma; Z)_{(p)} = \prod_{P_j \in S} \hat{H}^*(N(P_j); Z)_{(p)}$, where S is the set of all conjugacy classes of Z/p of Γ . Therefore, the p -period of Γ is the least common multiple of the p -periods of $N_\Gamma(P_i)$.

- 1) Lower bound. Let $|N_\Gamma(P_i)/C_\Gamma(P_i)| = d_i$, $\langle x \rangle = P_i$. There exists $y \in \Gamma$, such that $xyx^{-1} = x^r$, $r^{d_i} = 1 \pmod{p}$. Let $H = \langle x, y \rangle$ be a subgroup of Γ generated by elements x and y . Then the p -period of H is $2d_i$ by Lemma 1.1, i.e., the p -period of $N_\Gamma(P_i)$ is a multiple of $2d_i$.
- 2) Upper bound. Let $pr : \Gamma \rightarrow M$ be a projection onto the finite quotient M whose a p -Sylow subgroup is elementary abelian, and $pr_i : N_\Gamma(P_i) \rightarrow M_i$ be the restriction map of pr , where M_i is the image of pr_i . Then $M_i = Im N_\Gamma(P_i) = N_{M_i}(P_i)$ normalizes P_i (P_i also denotes the image of P_i), the group A_p of automorphisms of S_p induced by inner automorphisms of M_i maps P_i to itself.

Let $u \in H^2(S_p; Z) = \text{Hom}(P_i \times Z/p \times \dots \times Z/p, C^*)$ be a cohomology element such that $u(x) \neq 1$ and $u(y) = 1$ if $\langle x \rangle = P_i$, $\langle y \rangle = Z/p$, where C^* is the multiple group of nonzero complex numbers. Then $\text{Res}(u) \neq 0$ in $H^2(P_i; Z)$. Now we claim that $u^{d_i} \in H^{2d_i}(S_p; Z)$ is a stable element for S_p in M_i . In fact, $d_i = |N_{M_i}(P_i)/C_{M_i}(P_i)|$ by Lemma 1.2, and A_p fixes the element $u^{d_i} \in H^{2d_i}(S_p; Z)$ since $N_{M_i}(P_i)/C_{M_i}(P_i)$ fixes the element u^{d_i} . By Lemma 1.4 [9], u^{d_i} is a stable element for S_p in M_i , i.e., there exists an element $v \in H^{2d_i}(M_i; Z)$ such that $\text{Re } s_{P_i}^{M_i}(v) = \text{Re } s_{P_i}^{S_p}(u^{d_i}) = [\text{Re } s_{P_i}^{S_p}(u)]^{d_i} \neq 0$. If we apply the canonical homomorphism g^* from ordinary cohomology to Farrell cohomology [3, p. 278] we have $\text{Re } s_{P_i}^{M_i}(g^*(v)) = \text{Re } s_{P_i}^{S_p}(g^*(u^{d_i})) = \text{Re } s_{P_i}^{S_p}(g^*(u))^{d_i} \neq 0$, i.e., there exists an element $pr_i^* g^*(v) \in \hat{H}^{2d_i}(N_\Gamma(P_i); Z)$ such that $\text{Re } s_{P_i}^{N(P_i)}(pr_i^* g^*(v)) \neq 0$ in $\hat{H}^{2d_i}(P_i; Z)$, by Brown-Venkov theorem [2] and the fact that $N_\Gamma(P_i)$ has only one order p subgroup, the p -period of

$N_\Gamma(P_i)$ divides $2d_i$. See following diagram.

$$\begin{array}{ccccc}
 \hat{H}^{2d_i}(N_\Gamma(P_i); Z) & \xrightarrow{\text{Res}} & & & \hat{H}^{2d_i}(P_i; Z) \\
 \uparrow \text{pri}^* & & & & \uparrow \parallel \\
 \hat{H}^{2d_i}(M_i; Z) & \xrightarrow{\text{Res}} & \hat{H}^{2d_i}(S_p; Z) & \xrightarrow{\text{Res}} & \hat{H}^{2d_i}(P_i; Z) \\
 \uparrow \parallel g^* & & \uparrow \parallel g^* & & \uparrow \parallel g^* \\
 H^{2d_i}(M_i; Z) & \xrightarrow{\text{Res}} & H^{2d_i}(S_p; Z) & \xrightarrow{\text{Res}} & H^{2d_i}(P_i; Z) \blacksquare
 \end{array}$$

Proof of Theorem 2: is basically a similar argument except for the upper bound part. In fact, if Γ has a finite p -periodic quotient M with torsion free kernel, then Γ is p -periodic and the p -period of Γ divides the p -period of M . This is because the inflation map $\hat{H}^*(M) \rightarrow \hat{H}^*(\Gamma)$ maps an invertible element of $\hat{H}^*(M)$ to an invertible element of $\hat{H}^*(\Gamma)$. Using Swan Theorem, we obtain that the p -period of $N_\Gamma(P_i)$ divides the p -period of M_i , which is $2|N_{M_i}(Z/p^n)/C_{M_i}(Z/p^n)|$. Also, by Lemma 1.3, the number $2|N_{M_i}(Z/p^n)/C_{M_i}(Z/p^n)|$ divides $2|N_{M_i}(P_i)/C_{M_i}(P_i)| = 2|N_\Gamma(P_i)/C_\Gamma(P_i)|$. ■

2. An example

Lemma 1.3, Lemma 1.1 and Swan Theorem imply that the equality $|N(S_p)/C(S_p)| = |N(Z/p)/C(Z/p)|$ holds in the case of a finite group G whose a p -Sylow subgroup is cyclic, here Z/p is the order p subgroup of S_p . Therefore, Theorems 1 and 2 are generalizations of Swan Theorem.

In the case of a group Γ of finite vcd , in general, $|N(S_p)/C(S_p)| \neq |N(Z/p)/C(Z/p)|$ even if all maximal p -subgroups S_p of Γ are cyclic. For example, let $\Gamma^* = \langle x, y | x^{p^2} = 1, yxy^{-1} = x^{p+1} \rangle$, and d is the minimal positive integer such that $(p+1)^d \equiv 1 \pmod{p^2}$. Then $|N(\langle x \rangle)/C(\langle x \rangle)| = d = p$, but $|N(\langle x^p \rangle)/C(\langle x^p \rangle)| = 1$. A similar argument to Lemma 1.1 shows the p -period of Γ^* above equals $2p$. This trivial example shows that the p -period of an infinite group Γ can not be only described in the form $2LCM\{|N(Z/p)/C(Z/p)|\}$ in general.

The example Γ^* above could lead us to think that the p -period of a p -periodic group Γ equals $2LCM\{|N(C(p))/C(C(p))|\}$, where $C(p)$ ranges over all conjugacy classes of maximal p -cyclic subgroups of Γ . Recall in the case of a finite group G , Swan Theorem can be also stated in the different form: the p -period of G equals $2|N(C(p))/C(C(p))|$ (including the case $p = 2$), where $C(p)$ is a maximal p -cyclic subgroup of G .

Unfortunately, the next example shows that this is not true.

Example. Let $\Gamma_{n,m}$ denote the congruence subgroup of $SL(n, Z)$ of level m , i.e., the kernel of the surjective homomorphism $r_m : SL(n, Z) \rightarrow SL(n, Z/m)$ induced by the reduction mod(m) (m may not be prime). It is well-known that the group $\Gamma_{n,m}$ is always torsion free when $n \geq 1$ and $m \geq 3$. A result of Charney [4] states that the group $\Gamma_{n,p}$ is cohomology stable with $Z/2$ coefficient for any odd prime p . Define $\Gamma_p = \lim_n \Gamma_{n,p}$, then $H^i(\Gamma_{n,p}; Z/2) = H^i(\Gamma_p; Z/2)$ for $n \geq 2i + 5$.

Let $GL(Z)$ be the infinite general linear group of Z and $w_i \in H^i(GL(Z); Z/2)$ the i -th Stiefel-Whitney class of the inclusion $GL(Z) \rightarrow GL(R)$ for $i \geq 1$. We still denote by w_i the image of w_i under the restriction $H^i(GL(Z); Z/2) \rightarrow H^i(SL(Z); Z/2) \rightarrow H^i(\Gamma_m; Z/2)$.

The calculation in [1] by Arlettaz gives following results: for any odd prime p

- a) $w_1(\Gamma_p) = 0$
- b) $w_2(\Gamma_p) \neq 0$
- c) $w_3(\Gamma_p) = 0$ if and only if $p = 7 \pmod{8}$.

Also, we know from Wu formula for the Steenrod square $Sq^1(w_2) = w_1w_2 + w_0w_3 = w_3$ in $H^3(\Gamma_p; Z/2)$. Again, denote by w_i the image of w_i under the restriction $H^i(\Gamma_5; Z/2) \rightarrow H^i(\Gamma_{11,5})$. Combining both results of Charney and Arlettaz above, we have $w_1 = 0, w_2 \neq 0$ and $Sq^1(w_2) = w_3 \neq 0$ in $H^*(\Gamma_{11,5}; Z/2)$ (in fact, these are all true for $H^*(\Gamma_{n,5}; Z/2)$ as long as $n \geq 11$.)

Let Γ_0 denote the group of the extension $1 \rightarrow Z/2 \rightarrow \Gamma_0 \rightarrow \Gamma_{11,5} \rightarrow 1$ which corresponds to the non-trivial cohomology element $w_2 \in H^2(\Gamma_{11,5}; Z/2)$. Obviously, the group Γ_0 contains only one 2-subgroup $Z/2$, and the extension is central. Next, we check that the group Γ_0 is of finite vcd , then show that the 2-period of Γ_0 is greater than 2.

Consider the following commutative diagram, where all maps R_1, R_2, R_3 and R_4 are restriction maps.

$$\begin{array}{ccc}
 H^2(\Gamma_{11,4}; Z/2) & \xrightarrow{R_3} & H^2(\Gamma_{11,20}; Z/2) \\
 \uparrow R_1 & & \uparrow R_2 \\
 H^2(SL(11, Z); Z/2) & \xrightarrow{R_4} & H^2(\Gamma_{11,5}; Z/2)
 \end{array}$$

In fact, the map $R_1 = 0$ is a special case of the result by Millson [7, p. 85] which states that for any $n \geq 3$ the map $r^* : H^2(SL(n, Z/4); Z/2) \rightarrow H^2(SL(n, Z), Z/2)$ induced by the reduction mod(4) is an isomorphism.

Thus, we obtain the nontrivial second Stiefel-Whitney class w_2 in $H^2(\Gamma_{11,5}; Z/2)$, but the restriction of w_2 into the cohomology of the finite index subgroup $H^2(\Gamma_{11,20}; Z/2)$ is 0. This actually proves that the group Γ_0 is finite vcd and the $vcd(\Gamma_0) = cd(\Gamma_{11,20}) = vcd(SL(11, Z)) = 55$ [3, p. 229].

In order to find a lower bound on the 2-period of Γ_0 , consider two spectral sequences as follows:

1. The Lyndon-Hochschild-Serre spectral sequence of the group extension $1 \rightarrow Z/2 \rightarrow \Gamma_0 \rightarrow \Gamma_{11,5} \rightarrow 1$ with $Z/2$ coefficient. This takes the form $E_2^{i,j} = H^i(\Gamma_{11,5}; H^j(Z/2; Z/2)) \Rightarrow H^{i+j}(\Gamma_0; Z/2)$.
2. The Farrell cohomology spectral sequence [2] of the group extension $1 \rightarrow Z/2 \rightarrow \Gamma_0 \rightarrow \Gamma_{11,5} \rightarrow 1$ with $Z/2$ coefficient. This takes the form $E_2^{i,j} = H^i(\Gamma_{11,5}; \hat{H}^j(Z/2; Z/2)) \Rightarrow \hat{H}^{i+j}(\Gamma_0; Z/2)$.

Let $u \in H^1(Z/2; Z/2)$ be the generator of the cohomology ring $H^*(Z/2; Z/2) = F_2[u]$, and $d_2(u) = w_2 \in H^2(\Gamma_0; Z/2)$ be the second Stiefel-Whitney class corresponding to the extension $1 \rightarrow Z/2 \rightarrow \Gamma_0 \rightarrow \Gamma_{11,5} \rightarrow 1$. Then u is transgressive, $d_2(u) = \tau(u) = w_2$, where τ is the transgression. The element $u^2 = Sq^1(u)$ is also transgressive [8, p. 81], and $d_3(u^2) = \tau(u^2) = \tau(Sq^1(u)) = Sq^1(\tau(u)) = Sq^1(w_2) = w_3 \neq 0$ in E_3 because $H^1(\Gamma_{11,5}; Z/2)$ is trivial.

Consider a commutative diagram involving in both spectral sequences as follows:

$$\begin{array}{ccc}
 H^0(\Gamma_{11,5}; \hat{H}^2(Z/2; Z/2)) & \xrightarrow{d_3} & H^3(\Gamma_{11,5}; \hat{H}^0(Z/2; Z/2)) \\
 \uparrow \parallel g^* & & \uparrow \parallel g^* \\
 H^0(\Gamma_{11,5}; H^2(Z/2; Z/2)) & \xrightarrow{d_3} & H^3(\Gamma_{11,5}; H^0(Z/2; Z/2))
 \end{array}$$

The nontriviality of d_3 in the second row implies the nontriviality of d_3 in the first row. This shows $\text{Res}: \hat{H}^2(\Gamma_0; Z/2) \rightarrow \hat{H}^2(Z/2; Z/2)$ is trivial since the map Res factors through $E_\infty^{0,2} = 0$. Therefore, there is no invertible element in $\hat{H}^2(\Gamma_0; Z/2)$. By the fact that the reduced map $\hat{H}^2(\Gamma_0; Z)_{(2)} \rightarrow \hat{H}^2(\Gamma_0; Z/2)$ is ring homomorphism, there is no invertible element in $\hat{H}^2(\Gamma_0; Z)_{(2)}$, i.e., the 2-period of Γ_0 is greater than 2. We have proved our Theorem 3.

3. The p -period of the mapping class group Γ_g

The p -periodicity of the mapping class group is studied in a different paper of the author [11]. As an application of the theorem 1, we obtain the p -period of a p -periodic mapping class group Γ_g when $g < p(p-1)/2$.

Recall that the mapping class group Γ_g is defined to be the group of path components of orientation preserving diffeomorphisms of the closed orientable surface S_g of genus $g > 1$. Next, we define a set $B_{g,p}$ for surface S_g and a prime p .

Definition. For p odd, let $2g - 2 = mp - i, 0 \leq i \leq p - 1$.

$$B_{g,p} = \{i, i + p, i + 2p, \dots \dots i + ([2g/(p - 1)] - m)p\} \text{ if } i \neq 1.$$

$$B_{g,p} = \{1 + p, 1 + 2p, \dots \dots 1 + ([2g/(p - 1)] - m)p\} \text{ if } i = 1.$$

And for $p = 2$,

$$B_{g,2} = \{0, 4, 8, \dots \dots 2g + 2\} \text{ if } g \text{ is odd.}$$

$$B_{g,2} = \{2, 6, 10, \dots \dots 2g + 2\} \text{ if } g \text{ is even.}$$

Remarks.

1. The notation $[-]$ here means the integer part.
 In case $i \neq 1, 2g/(p - 1) < m$, define $B_{g,p} = \emptyset$.
 In case $i = 1, 2g/(p - 1) < m + 1$, define $B_{g,p} = \emptyset$.
2. It is proved in [11] that the set $B_{g,p}$ is exactly the set of all possible number of fixed points when an order p diffeomorphism acts on the surface S_g .

Lemma 3.1. *For the mapping class group Γ_g , there is a formula $LCM\{|N(\langle x \rangle)/C(\langle x \rangle)|\} = LCM\{\gcd(p - 1, b_i)\}$, where $\langle x \rangle$ ranges over all conjugacy classes of Z/p in Γ_g , b_i ranges over all $b_i \in B_{g,p}$.*

Proof: 1) Assume $|N(\langle x \rangle)/C(\langle x \rangle)| = d$. Then there exists an integer r such that $x \approx x^r \approx \dots \dots \approx x^{r^{d-1}}$ (\approx means "is conjugate to" in Γ_g) so that d is the minimal positive integer satisfying $r^d = 1 \pmod{p}$. The d divides $p - 1$ obviously. Let b be the number of fixed points of the x action on S_g , $\sigma(x) = (\beta_1, \beta_2, \dots \dots \beta_b)$ the fixed point datum, where $\beta_i \in Z/p - \{0\}$ (cf. [10]).

Let us define a permutation r^* on the ordered b_i -tuple $(\beta_1, \beta_2, \dots, \beta_{b_i})$. Set $r^*(\beta_1, \beta_2, \dots \dots \beta_{b_i}) = (r\beta_1, r\beta_2, \dots \dots r\beta_{b_i})$, $(r^*)^2 = (r^2)^* \dots \dots (r^*)^{d-1} = (r^{d-1})^*$. It is well-defined since $\sigma(x) = \sigma(x^{r^2}) = \dots \dots = \sigma(x^{r^{d-1}})$ as an unordered b -tuples [12]. We can decompose $r^* = (\beta_{i_1}, \beta_{i_2}, \dots \dots \beta_{i_s})(\beta_{j_1}, \beta_{j_2}, \dots \dots \beta_{j_t}) \dots \dots (\beta_{k_1}, \beta_{k_2}, \dots \dots \beta_{k_u})$, a product of cyclic permutations. Notice that permutations $r^*, (r^*)^2, \dots \dots (r^*)^{d-1}$ do not have fixed points. Otherwise, there exists β_i such that $rj\beta_i = \beta_i \pmod{p}$, $1 \leq j \leq d - 1$. This forces $rj = 1 \pmod{p}$, a contradiction. But, of course, $(r^*)^d = (r^d)^* = \text{Id}$. These imply

$s = t = \dots = u = d$, i.e., the number $|N(\langle x \rangle)/C(\langle x \rangle)| = d$ divides the number b_i of fixed points of the x action on the surface S_g . We have showed that $LCM\{|N(\langle x \rangle)/C(\langle x \rangle)|\}$ divides $LCM\{\gcd(p-1, b_i)\}$, where $\langle x \rangle$ ranges over all conjugacy classes of Z/p in Γ_g , b_i ranges over all $b_i \in B_{g,p}$.

2) Conversely, assume $\gcd(p-1, b_i) = d$. Then there is a mod(p) integer r so that d is a minimal positive integer satisfying $r^d = 1 \pmod{p}$.

Case 1. $b_i \neq 0$. If $d \neq 1$, then $r \neq 1$. Consider the unordered b_i -tuples $\sigma = (1, r, r^2, \dots, r^{d-1}, 1, r, r^2, \dots, r^{d-1}, \dots, 1, r, r^2, \dots, r^{d-1})$. Since $(b_i/d)(1+r+r^2+\dots+r^{d-1}) = 0 \pmod{p}$. There exists an element $x \in \Gamma_g$, $x^p = 1$, and the it's representative fixed point datum $\sigma(x)$ is σ , i.e., the unordered b_i -tuples σ can be realized as a fixed point datum of an order p element in Γ_g [6]. Obviously, $\sigma(x) = \sigma(x^r) = \sigma(x^{r^2}) = \dots = \sigma(x^{r^{d-1}})$ or $x \approx x^r \approx x^{r^2} \approx \dots \approx x^{r^{d-1}}$ in Γ_g . This implies that the number d divides the order $|N(\langle x \rangle)/C(\langle x \rangle)|$. If $\gcd(p-1, b_i) = d = 1$, for any order p element x in Γ_g with the number of fixed points b_i , obviously 1 divides $|N(\langle x \rangle)/C(\langle x \rangle)|$.

Case 2. $b_i = 0$. On the one hand, we have $\gcd(p-1, b_i) = p-1$. On the other hand, the x acts on S_g freely. All order p free actions are conjugate by [5], this implies $|N(\langle x \rangle)/C(\langle x \rangle)| = p-1$.

So, $LCM\{\gcd(p-1, b_i)\}$ divides $LCM\{|N(\langle x \rangle)/C(\langle x \rangle)|\}$. ■

Proof of Theorem 4: Let $\mu : \Gamma_g \rightarrow Sp(2g, Z)$ be the canonical homology representation and $p : Sp(2g, Z) \rightarrow Sp(2g, F_q)$ be the reduction map. Here q can be chosen a primitive root of mod(p) such that $q \geq 3$, and q^{p-1} is not congruent to 1 mod(p^2) (by the Dirichlet theorem).

Now $\text{Ker}(p\mu) = N$ is a torsion free, normal, finite index subgroup of Γ_g and a p -Sylow subgroup of the finite quotient $\Gamma_g/N = Sp(2g, F_q)$ is elementary abelian if $2g < p(p-1)$. Then we can use Theorem 1 and Lemma 3.1 to finish the proof. ■

A list of the p -period of a p -periodic mapping class group Γ_g can be also found in the Appendix C of the author's thesis [12].

Acknowledgement. The most parts of this paper are from the Ohio State University thesis [13] which is written by the author under the direction of Professor H. Glover and Professor G. Mislin. The author wishes to express the most sincere thanks to Professor H. Glover and Professor G. Mislin for their excellent guidance and inspiration.

References

1. ARLETTAZ, D., Sur les classes de Stiefel-Whitney des sous-groupes

- de congruence, *C.R. Acad. Sci. Paris Ser. I Math.* **303** (1986), 571-574.
2. BROWN, K.S., Groups of virtually finite dimension, Proc. of Sept, 1977, Durham conference on homological and combinatorial techniques in group theory, 27-70.
 3. BROWN, K.S., "*Cohomology of groups*," Graduate Texts in Math. **87**, Springer-Verlag, 1982.
 4. CHARNEY, R., On the problem of homology stability for congruence subgroups, *Comm. in Algebra* **12** (1984), 2081-2123.
 5. EDMONDS, A., Surface symmetry I, *Mich. Math. J.* **29** (1982), 171-183.
 6. EWING, J., "*Automorphisms of surfaces and class numbers: An illustration of the G-index theorem*," London Math. Soc. Lecture note series **86**, 1983, pp. 120-127.
 7. MILLSON, J., Real vector bundles with discrete structure group, *Topology* **18** (1979), 83-89.
 8. MOSHER, R. AND TANGORA, M., "*Cohomology operations and Applications in homotopy theory*," Harper and Row, Publishers.
 9. SWAN, R., The p -period of a finite group, *Illinois J. Math.* **4** (1960), 341-346.
 10. SYMONDS, P., The cohomology representation of an action of C_p on a surface, *Trans. Amer. Math. Soc.* **306** (1988), 389-400.
 11. XIA, Y., The p -periodicity of the mapping class group and the estimate of its p -period, *Proc. of AMS*, to appear.
 12. XIA, Y., Farrell-Tate cohomology of the mapping class group, Thesis, The Ohio State University.

Department of Mathematics
The Ohio State University
Columbus, Ohio 43210-1174
U.S.A.

Primera versió rebuda el 21 de Juny de 1991,
darrera versió rebuda el 13 de Febrer de 1992