

## A variant of the Smale's model for flocks formation

J. I. Díaz

**Abstract.** Motivated by a recent lecture of S. Smale, we propose here a variant of the model considered by F. Cucker and S. Smale for the formation of uniform flocks allowing to prove that the formation of an uniform structure take place in a finite time (and not merely when  $t$  converges to infinity).

### Una variación sobre un modelo de Smale para la formación de bandadas

**Resumen.** Motivado por una reciente conferencia de S. Smale, proponemos aquí una variación del modelo considerado por F. Cucker y S. Smale que permite mostrar la formación de bandadas con un movimiento sincronizado en un tiempo finito (y no meramente cuando  $t$  converge a infinito).

## 1 Introduction

This note is motivated by a part of a lecture by S. Smale ([7]) in which he reported some results on his previous work [2], in collaboration with F. Cucker, on the formation of flocks by birds (or fish), according a model which proposed by Vicsek et al. [8]. The motivation comes from the observation that under some conditions (for example on the structure of its dynamics) the state of the flock converges to one in which all birds fly with the same velocity. The goal of the above mentioned papers was to provide some justification of this observation.

We propose here a slight variation of the simplified model for the case of two birds

$$P(\alpha, \beta) \begin{cases} \dot{x} = v, \\ \dot{v} = -\beta \frac{v^{(2-\frac{1}{\beta})}}{(1+x)^\alpha}, \\ x(0) = x_0, \\ v(0) = v_0, \end{cases} \quad (1)$$

where  $\alpha, \beta \in (0, 1]$ .

The case  $\beta = 1$  and  $\alpha > 1$  was considered in [2, Section 4] (see also the general results, including the case  $\alpha \in (0, 1)$  in [2, Section 3]). They prove that the uniform folk is formed asymptotically when  $t \rightarrow +\infty$ .

Our main goal is to prove the finite (continuous) time convergence to a flock (we recall that the convergence in discrete time was obtained in [2, Section 5] for a discrete version of a general model, including (1) with  $\beta = 1$  as a special case).

---

Presentado por Manuel López Pellicer.

Recibido: 11 de julio de 2007. Aceptado: 10 de octubre de 2007.

Palabras clave / Keywords: Flocks, finite time convergence to equilibrium, nonlinear ordinary differential systems

Mathematics Subject Classifications: 34C99, 34A34.

© 2007 Real Academia de Ciencias, España.

**Theorem 1** Assume  $\alpha \in (0, 1)$ ,  $v_0 > 0$  and  $x_0 > 0$ . Then there exists  $T_f = T_f(x_0, v_0) \in (0, +\infty]$  such that, when  $t \rightarrow T_f$ ,  $v(t) \rightarrow 0$  and

$$x(t) \rightarrow x_\infty := \left( (1 - \alpha)v_0^{1/\beta} + (1 + x_0)^{1-\alpha} \right)^{\frac{1}{1-\alpha}} - 1. \quad (2)$$

Moreover,  $x(t)$  is strictly increasing on  $[0, T_f)$  and we have that  $T_f = +\infty$  if  $\beta = 1$  and  $T_f < +\infty$  if  $\beta \in (0, 1)$ . In the case,  $\beta \in (0, 1)$ , the solution satisfies that

$$x(t) = x_\infty \quad \text{and} \quad v(t) = 0 \quad \text{for any } t \in [T_f, +\infty).$$

IDEA OF THE PROOF. We start by recalling that the existence of a local solution comes from the continuity of the right hand side of both equations. The uniqueness of solution can be obtained (even for the case  $\alpha \in (0, 1)$  and  $\beta \in (0, 1)$ ) thanks to the monotonicity of the application  $v \rightarrow v^{(2-\frac{1}{\beta})}$  and by using that the application  $x \rightarrow \frac{1}{(1+x)^\alpha}$  is locally Lipschitz continuous (recall that  $v_0 > 0$  and  $x_0 > 0$ ). We point out that, if we define

$$V = v^{1/\beta}$$

then the system can be reformulated as

$$\tilde{P}(\alpha, \beta) \begin{cases} \dot{x} = V^\beta, \\ \dot{V} = -\frac{V^\beta}{(1+x)^\alpha}, \\ x(0) = x_0, \\ V(0) = V_0 := v_0^{1/\beta}. \end{cases} \quad (3)$$

As a matter of fact, we shall prove our conclusion by using this equivalent formulation. Since the system  $\tilde{P}(\alpha, \beta)$  is autonomous we can eliminate the time on each orbit and write, equivalently the orbit as solution of the scalar equation

$$\frac{dx}{dV}(x) = -(1+x)^\alpha. \quad (4)$$

Thus, we have that

$$V(t) - V_0 = \frac{1}{(1-\alpha)} [(1+x_0)^{1-\alpha} - (1+x(t))^{1-\alpha}]. \quad (5)$$

From here and the fact that  $v_0 > 0$  and  $x_0 > 0$  we deduce that the pair  $(x(t), V(t))$  remains in a bounded set of  $\{(x, V) \in \mathbb{R}^2, x \geq 0, V \geq 0\}$  when  $t \in [0, +\infty)$  and since  $\dot{x} \geq 0$  and  $\dot{V} \leq 0$  we get that, necessarily,  $x(t) \rightarrow x_\infty$ , for some  $x_\infty \geq x_0$ , and  $v(t) \rightarrow 0$  when  $t \rightarrow +\infty$ . Moreover, from (5) we get that  $x_\infty$  is given by (2). Assume now that  $\beta \in (0, 1)$ . Let us prove that the limits are attained in a finite time. By introducing

$$\hat{x}(t) = 1 + x(t)$$

and using the first equation of (3) we get that

$$\frac{d\hat{x}}{dt}(t) = \frac{1}{(1-\alpha)} (\hat{x}_\infty^{1-\alpha} - \hat{x}(t)^{1-\alpha})^\beta,$$

where  $\hat{x}_\infty = 1 + x_\infty$ . In particular, for any  $t \geq 0$

$$\int_{\hat{x}_0}^{\hat{x}(t)} \frac{ds}{(\hat{x}_\infty^{1-\alpha} - s^{1-\alpha})^\beta} = t,$$

where  $\hat{x}_0 = x_0 + 1$ . Thus,  $x(t) \rightarrow x_\infty$  when  $t \rightarrow T_f$ , for a finite time  $T_f > 0$ , if and only if the following improper integral is convergent

$$\int_{\hat{x}_0}^{\hat{x}_\infty} \frac{ds}{(\hat{x}_\infty^{1-\alpha} - s^{1-\alpha})^\beta} < +\infty.$$

But this can be easily checked (make the change of variables  $u = \hat{x}_\infty^{1-\alpha} - s^{1-\alpha}$ ) thanks to the assumption  $\beta \in (0, 1)$ .  $\square$

**Remark 1** A very rich situation arises when  $\alpha \in (0, 1)$  and  $x_0 = 0$  (even for  $\beta = 1$ ), which seems to be unadvertised in [2]. As we deduce from the auxiliary equation

$$\frac{d\hat{x}}{dV}(x) = -\hat{x}(V)^\alpha$$

now for  $\hat{x}(V_0) = 0$  we get a continuum of solutions (which may remain satisfying  $\hat{x}(V) = 0$  on the interval  $[V_s, V_0]$  for an arbitrary  $V_s \in (0, V_0]$  and  $\hat{x}(V) > 0$  and  $\hat{x}(V) \rightarrow \hat{x}_\infty$  when  $V \rightarrow 0$ ). The above convergence  $\hat{x}(V) \rightarrow \hat{x}_\infty$  when  $V \rightarrow 0$  takes place in a infinite (respectively a finite time) if  $\beta = 1$  (respectively  $\beta \in (0, 1)$ ). A curious fact, considered in [1], is that this continuum of solutions still exists for equations of the form

$$\frac{d\hat{x}}{dV}(x) = -\hat{x}(V)^\alpha + f(\hat{x}(V))$$

when

$$|f(s)| \leq cs^{\frac{\alpha}{1-\alpha}}$$

for some  $c > 0$  small enough.

**Remark 2** As mentioned in [2], model  $P(\alpha, \beta)$  (even for  $\beta = 1$ ) is only a simplification of a vectorial model of the type

$$\mathbf{P}(L) \begin{cases} \dot{\mathbf{x}} = \mathbf{v}, \\ \dot{\mathbf{v}} = -L(\mathbf{x})(\mathbf{v}), \\ \mathbf{x}(0) = \mathbf{x}_0, \\ \mathbf{v}(0) = \mathbf{v}_0, \end{cases} \quad (6)$$

where  $\mathbf{x}, \mathbf{v} \in \mathbb{R}^{3k}$  and  $L(\mathbf{x})$  denotes a matrix, called as the Laplacian of the ‘‘adjacency matrix’’  $A$

$$A(\mathbf{x}) = (a_{ij}) = (\eta(\|\mathbf{x}_i - \mathbf{x}_j\|^2))$$

where  $\mathbf{x}_i \in \mathbb{R}^3$ ,  $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k) \in \mathbb{R}^{3k}$  and

$$\eta(y) = \frac{K}{(\sigma^2 + y)^\alpha}.$$

In fact, in the case of two birds following parallel lines (the scalar case) we must understand that  $x = x_1 - x_2$  and  $v = v_1 - v_2$ . By using some energy arguments (as in [1]), it seems possible to extend Theorem 1 to some suitable systems of the type  $\mathbf{P}(\alpha, \beta)$  when the matrix  $L(\mathbf{x})$  depends also on  $\mathbf{v}$  in a suitable way.

**Remark 3** Discrete models corresponding to a time discretization of  $\mathbf{P}(L)$  are relevant in learning and language evolution ([3, 6, 7]). It is not difficult to extend the conclusions of Theorem 1 to the discrete model associated to problem  $P(\alpha, \beta)$  (see, some related results in [4] and its references).

**Acknowledgement.** Partially supported by the project MTM2005-03463 of the DGISGPI (Spain) and CCG06-UCM/ESP-1110 of the DGUIC of the CAM and the UCM.

## References

- [1] Antontsev, S. N., Díaz, J. I. and Shmarev, S. I., (2002). *Energy Methods for Free Boundary Problems: Applications to Non-linear PDEs and Fluid Mechanics*, Birkhäuser, Boston, Progress in Nonlinear Differential Equations and Their Applications, **48**.
- [2] Cucker, F. and Smale, S., (2007). Emergent behavior in Flock, *IEEE Transactions on automatic control*, **52**, 5, 852–862.
- [3] Cucker, F., Smale, S. and Zhou, D.-X., (2004). Modeling language evolution, *Found. Comput. Math.*, **4**, 315–343.
- [4] Díaz, J. I. *Energy methods for free boundary problems: new results and some remarks on numerical algorithms*, Serie électronique de la SMAI, Esaim Proc.
- [5] Flierl, G., Grünbaum, D., Levin, S. and Olson, D., (1999). From individuals to aggregations: the interplay between behavior and physics, *J. Theor. Biol.*, **196**, 397–454.
- [6] Ke, J., Minett, J., Au, C.-P. and Wang, W.S.-Y., (2002). Self-organization and selection in the emergence of vocabulary, *Complexity*, **7**, 41–54.
- [7] Smale, S. (2007). *The changing directions of Mathematics*, Lecture at the International Symposium “The frontiers of Mathematics”, Fundación Ramón Areces, Madrid, June 7-8, 2007. Organized by the CSIC and the RAC.
- [8] Vicsek, T., Czirók, A., Ben-Jacob, E. and Shochet, O., (1995). Novel type of phase transition in a system of self-driven particles. *Phys. Rev. Letters*, **75**, 1226–1229.

**J. I. Díaz**

Dpto. de Matemática Aplicada Facultad de Matemáticas

Universidad Complutense de Madrid

28040 Madrid Spain

diaz.racefyn@insde.es