

Remarks on the Symmetries of Planar Fronts

F. AICARDI

ABSTRACT. A front is the projection on the plane of a legendrian immersion of a circle in the space of the contact elements of that plane. I analyse the symmetries of a generic front with respect to the group generated by the involutions reversing the orientation of the plane, the orientation of the preimage circle and the coorientation of the contact plane.

1. GENERIC FRONTS

A *planar front* is the projection to \mathbf{R}^2 (with coordinates x, y) of a legendrian curve. A *legendrian curve* is the image of a C^1 -immersion of S^1 in the space M^3 (with coordinates $x, y, \phi(\text{mod } 2\pi)$ for cooriented fronts, $(\text{mod } \pi)$ for noncooriented fronts) of the contact elements of the plane, with its natural contact structure

$$(\cos\phi)dx + (\sin\phi)dy = 0 \quad (1)$$

We call such an immersion of S^1 into M^3 an *L-immersion*

The front is *cooriented*, if the contact element is cooriented, i.e. if a choice of one of two halfplanes into which it divides the tangent plane is made. We shall consider now the cooriented fronts, and M^3 will be the space of the cooriented contact elements.

The front is *oriented* if the preimage circle S^1 is oriented.

A *generic front* may have as singularities only the ordinary double points and the semicubical cusp points.

Examples. See fig.1.




Fig. 1

Oriented legendrian curves and their oriented and cooriented fronts

To any oriented legendrian curve in the space of cooriented contact elements of the plane one associates two integer numbers: the *index* and the *Maslov index* (see [1]). Both these indices can be calculated in terms of the front of the legendrian curve.

Theorem 1. [1] *The index i of an oriented legendrian curve in the space M^3 is equal to the total angle (divided by 2π) of the rotation of the coorienting normal vector of its front when the point of the front makes a full turn along it.*

The Maslov index μ of a generic oriented legendrian curve is equal to the difference between the number μ^+ of positive cusps and the number μ^- of negative cusps of its front: $\mu = \mu^+ - \mu^-$.

A cusp of an oriented and cooriented front is called *positive* if the coorienting normal vector at the cusp point belongs to the halfplane bounded by the tangent line at the cusp and containing the cusp branch with the orientation going away from the cusp point.

Remarks (important) 1) The sign of the angle of rotation of the normal vector does not depend on the front coorientation, but depends on the orientation of the plane. Hence *the sign of the index does not depend on the front coorientation and changes when the orientations of the plane or of the front is changed.*

2) The sign of a cusp changes when one changes either the orientation or the coorientation of the front, but does not depend on the orientation of the plane. Hence *the Maslov index changes its sign when the orientation or the coorientation of the front is changed and it is independent of the orientation of the plane.*

Examples. See fig. 2




Fig. 2
Two fronts with their indices i and μ .

2. FRONT CLASSES AND SYMMETRY CLASSES

A *front class* is a class of L -immersions having generic fronts up to orientations preserving diffeomorphisms preserving the orientations of the plane and of the preimage circle.

Consider a group G acting on the space of L -immersions.

Definition 1. A front class $[\gamma]$ is called G -invariant if $[G\gamma] = [\gamma]$

Example. Let G_3 be the group of the euclidean motions in the plane. Any front class is G_3 -invariant.

Let T (T from Turn) ($T = S^1 \times S^1$) the group of the rotations of the plane and of the rotations of the preimage circle.

Remark. Every front class is invariant with respect to the group T .

Theorem 2. *If the front class $[\gamma]$ is invariant with respect to the symmetry group H , then there exist a front $\gamma_* \in [\gamma]$ so that $H\gamma_* \in T\gamma_*$, i.e. for any element h of H there exist an element τ in T so that*

$$h \gamma_* = \tau \gamma_*. \quad (2)$$

This theorem is stated in [2] with no proof for some special finite symmetry groups H .

We first consider the classification of oriented and cooriented fronts in the oriented plane up to the diffeomorphisms of the plane and of the preimage circle *preserving the orientations*. The front being cooriented, these diffeomorphisms preserve also the coorientation.

Let $\Omega_{i,\mu}$ be the space of immersions of the oriented circle into the space of the cooriented contact elements of the oriented plane with fixed index i and Maslov index μ , and $\mathcal{N}_{|i|,|\mu|}$ the corresponding space if only the absolute values of the indices are fixed. In this second space, consider the action of the following involutions:

Involution Σ : it reverses the orientation of the plane.

Involution σ : it reverses the orientation of the preimage circle.

Involution Θ : it reverses the coorientation of the front.

We represent an element of $\mathcal{N}_{|\lambda|,|\mu|}$ as the graph of the application of the circle into M^3 , i.e. as a curve $\tilde{\gamma}$ living in the space $S^1 \times M^3$ with coordinates

$$t(\text{mod}2\pi), \quad x(t), \quad y(t), \quad \phi(t)(\text{mod}2\pi).$$

The action of an element $\tau_{(\alpha,\theta)}$ of T on an L -immersion, i.e. the rotation of the preimage circle of an angle α and the rotation of an angle θ in the plane (x, y) is thus represented by the following transformation of $\tilde{\gamma}$ into $\tilde{\gamma}'$:

$$\begin{aligned} t' &= t + \alpha \\ x' &= (\cos\theta)x - (\sin\theta)y \\ y' &= (\cos\theta)y + (\sin\theta)x \\ \phi' &= \phi + \theta \end{aligned} \tag{3}$$

Moreover the involutions Σ , σ and Θ acting on the front classes can be represented with the following matrices acting on the immersions respectively:

involution Σ :

$$a = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \tag{4}$$

involution σ :

$$b = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{5}$$

involution Θ :

$$c = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{6}$$

These involutions generate a commutative group RO (Reversing Orientations) of order 8, whose elements coincide with their inverse ones. They are denoted by

$$I, a, b, c, ab, ac, bc, abc.$$

Let H be the symmetry group RO or a subgroup of RO . We want find the classes invariant under the action of the symmetry group H . Hence, using theorem 2, we search for a solution $(\tilde{\gamma}_H, \tau_i)$ of the system of equations

$$h_i \tilde{\gamma}_H = \tau_i \tilde{\gamma}_H \tag{7}$$

where h_i are the generators of the group H and τ_i are particular elements of the group T . If such a solution exists, then the front class containing γ_H will be the requested H -invariant class.









$i = 2 \quad \mu = 2$	$i = -2 \quad \mu = 2$	$i = -2 \quad \mu = -2$	$i = 2 \quad \mu = -2$
 [γ]	 [$a\gamma$]	 [$b\gamma$]	 [$c\gamma$]
 [$abc\gamma$]	 [$bc\gamma$]	 [$ac\gamma$]	 [$ab\gamma$]

Fig. 3
The eight asymmetric classes

Definition 2. We call *asymmetric class* a front class which is not invariant by any subgroup of RO different from the identity.

The action of RO on an asymmetric class of $\mathcal{N}_{|i|,|\mu|}$ transforms it into 8 different asymmetric classes, with different values of the signs of the nonzero indices.

Example. See fig. 3.

Definition 3. We call *simply symmetric class* a front class which is invariant only with respect to a subgroup of order two of RO .

Every element ρ of RO different from the identity I generates a subgroup of RO of order 2. We denote by $R_{(\rho)}$ this subgroup, by $[\gamma_\rho]$ the class invariant only with respect to $R_{(\rho)}$, by $\tilde{\gamma}_\rho$ the curve, solution of equation (7), and by γ_ρ its corresponding front. The action of RO on a simply symmetric class transform it into 4 different simply symmetric classes in $\mathcal{N}_{|i|,|\mu|}$.

Definition 4. We call *supersymmetric class* a front class which is invariant with respect to a subgroup of order four of RO .

Two elements ρ_1 and ρ_2 of RO different from the identity I generate a subgroup of RO of order 4 ($I, \rho_1, \rho_2, \rho_1\rho_2$). We denote by $R_{(\rho_1)(\rho_2)}$ this subgroup, by $[\gamma_{\rho_1,\rho_2}]$ its invariant class, by $\tilde{\gamma}_{\rho_1,\rho_2}$ the curve, solution of equation (7) and by γ_{ρ_1,ρ_2} its corresponding front. The action of RO on this class transforms it into 2 different supersymmetric classes in $\mathcal{N}_{|i|,|\mu|}$

Theorem 3. i) *There are no invariant classes under the action of the entire group RO .* ii) *The front classes invariant with respect to the subgroups $R_{(a)}, R_{(c)}, R_{(ab)}, R_{(ac)}, R_{(bc)}, R_{(abc)}, R_{(a)(bc)}, R_{(c)(ab)}, R_{(ab)(bc)}$ are not void. They are listed in the following table, where the solutions of eq. (7) for every subgroup are shown in the identities in the second column.*

Subgroup	Identities	Number of inv. cl.	Space of inv. cl.	Examples in figg.
$R_{(a)} = (I, a)$	$a\tilde{\gamma}_a = \tau_{\pi,0} \tilde{\gamma}_a$	4	$\mathcal{N}_{0, \mu }$	4, 5
$R_{(c)} = (I, c)$	$c\tilde{\gamma}_c = \tau_{\pi+\alpha,\alpha} \tilde{\gamma}_c (*)$	4	$\mathcal{N}_{ i ,0}$	6, 7
$R_{(ab)} = (I, ab)$	$ab\tilde{\gamma}_{ab} = \tau_{0,0} \tilde{\gamma}_{ab}$	4	$\mathcal{N}_{ i ,0}$	8, 9
$R_{(ac)} = (I, ac)$	$ac\tilde{\gamma}_{ac} = \tau_{\pi,0} \tilde{\gamma}_{ac}$	4	$\mathcal{N}_{0,0}$	10, 11
$R_{(bc)} = (I, bc)$	$bc\tilde{\gamma}_{bc} = \tau_{0,0} \tilde{\gamma}_{bc}$	4	$\mathcal{N}_{0, \mu }$	12, 13
$R_{(abc)} = (I, abc)$	$abc\tilde{\gamma}_{abc} = \tau_{0,0} \tilde{\gamma}_{abc}$	4	$\mathcal{N}_{ i , \mu }$	14, 15
$R_{(a),(bc)} = (I, a, bc, abc)$	$a\tilde{\gamma}_{a,bc} = \tau_{\pi,0} \tilde{\gamma}_{a,bc}$ $bc\tilde{\gamma}_{a,bc} = \tau_{0,0} \tilde{\gamma}_{a,bc}$	2	$\mathcal{N}_{0, \mu }$	16, 17
$R_{(c),(ab)} = (I, c, ab, abc)$	$c\tilde{\gamma}_{c,ab} = \tau_{\pi+\alpha,\alpha} \tilde{\gamma}_{c,ab} (*)$ $ab\tilde{\gamma}_{c,ab} = \tau_{0,0} \tilde{\gamma}_{c,ab}$	2	$\mathcal{N}_{ i ,0}$	18, 19
$R_{(ab)(bc)} = (I, ab, bc, ac)$	$ab\tilde{\gamma}_{ab,bc} = \tau_{\pi,0} \tilde{\gamma}_{ab,bc}$ $bc\tilde{\gamma}_{ab,bc} = \tau_{0,0} \tilde{\gamma}_{ab,bc}$	2	$\mathcal{N}_{0,0}$	20, 21

(*) $\alpha = (2n + 1)\frac{\pi}{p}$ (symmetry of order p even), or $\alpha = \frac{2n\pi}{p}$ (symmetry of order p odd).

Proof. Consider the subgroup $R_{(b)} = (I, b)$. The solution of the equation (7) for the symmetric front γ_b does not exist. According to theorem 2, there are not $R_{(b)}$ -invariant front classes. It follows that also the group RO cannot have invariant classes, because they would be symmetric with respect to this subgroup $R_{(b)}$.

The other subgroups of RO of order two have invariant classes. Indeed, according to theorem 2, the equation (7) has solutions in the forms shown in the table. Such solutions are unique, in the sense that

in every simply symmetric class $[\gamma_\rho]$ there exist an element γ_ρ which satisfies the corresponding identity shown in the second column. The simply symmetric fronts explicitly satisfy respectively:

the front γ_a satisfies:

$$\begin{aligned} -x(t) &= x(t + \pi) \\ y(t) &= y(t + \pi) \\ -\phi(t) &= \phi(t + \pi) \end{aligned} \tag{8}$$




Fig.4

A simply symmetric front γ_a with $i = 0$ and $\mu = 2$

the front γ_c satisfies:

$$\begin{aligned} -x(t) &= (\cos\alpha)x(t + \pi + \alpha) - (\sin\alpha)y(t + \pi + \alpha) \\ -y(t) &= (\cos\alpha)y(t + \pi + \alpha) + (\sin\alpha)x(t + \pi + \alpha) \\ \phi(t) &= \phi(t + \pi + \alpha) + \alpha \end{aligned} \tag{9}$$

where $\alpha = (2n + 1)\frac{\pi}{p}$ (symmetry of order p even), or $\alpha = \frac{2n\pi}{p}$ (symmetry of order p odd). An example of no rotation symmetry, i.e. the case $p = 1$ ($\alpha = 2\pi n$) is shown in fig. 6. An example of rotation symmetry ($p = 2$) is shown in fig.22




Fig.5
The four $R_{(a)}$ -invariant classes in $\mathcal{N}_{0,|\mu|}$




Fig.6
A simply symmetric front γ_c with $i = 2$ and $\mu = 0$




Fig.7

The four $R_{(c)}$ -invariant classes in $\mathcal{N}_{|i|,0}$
 the front γ_{ab} satisfies:

$$\begin{aligned}
 -x(-t) &= x(t) \\
 y(-t) &= y(t) \\
 -\phi(-t) &= \phi(t)
 \end{aligned}
 \tag{10}$$




Fig.8
A simply symmetric front γ_{ab} with $i = 1$ and $\mu = 0$

$i = 1 \quad \mu = 0$	$i = -1 \quad \mu = 0$
 $[\gamma_{ab}] = [ab\gamma_{ab}]$	 $[a\gamma_{ab}] = [b\gamma_{ab}]$
 $[c\gamma_{ab}] = [abc\gamma_{ab}]$	 $[ac\gamma_{ab}] = [bc\gamma_{ab}]$

Fig.9
The four $R_{(ab)}$ -invariant classes in $\mathcal{N}_{|i|,0}$

the front γ_{ac} satisfies:

$$\begin{aligned} x(t) &= x(t + \pi) \\ -y(t) &= y(t + \pi) \\ -\phi(t) &= \phi(t + \pi) \end{aligned} \tag{11}$$




Fig.10

A simply symmetric front γ_{ac} with $i = 0$ and $\mu = 0$




Fig.11

The four $R_{(ac)}$ -invariant classes in $\mathcal{N}_{0,0}$

the front γ_{bc} satisfies:

$$\begin{aligned} -x(-t) &= x(t) \\ -y(-t) &= y(t) \\ \phi(-t) &= \phi(t) \end{aligned} \tag{12}$$




Fig.12
A simply symmetric front γ_{bc} with $i = 0$ and $\mu = 2$





$i = 0 \quad \mu = 2$	$i = 0 \quad \mu = -2$
 $[\gamma_{bc}] = [bc\gamma_{bc}]$	 $[b\gamma_{bc}] = [c\gamma_{bc}]$
 $[a\gamma_{bc}] = [abc\gamma_{bc}]$	 $[ab\gamma_{bc}] = [ac\gamma_{bc}]$

Fig.13
The four $R_{(bc)}$ -invariant classes in $\mathcal{N}_{0,|\mu|}$

the front γ_{abc} satisfies:

$$\begin{aligned} x(-t) &= x(t) \\ -y(-t) &= y(t) \\ -\phi(-t) &= \phi(t) \end{aligned} \tag{13}$$




Fig.14

A simply symmetric front γ_{abc} with $i = 1$ and $\mu = 2$

$i = 1 \quad \mu = 2$	$i = 1 \quad \mu = -2$	$i = -1 \quad \mu = 2$	$i = -1 \quad \mu = -2$
$[\gamma_{abc}] = [abc\gamma_{abc}]$	$[c\gamma_{abc}] = [ab\gamma_{abc}]$	$[a\gamma_{abc}] = [bc\gamma_{abc}]$	$[b\gamma_{abc}] = [ac\gamma_{abc}]$

Fig.15

The four $R_{(abc)}$ -invariant classes in $\mathcal{N}_{|i|,|\mu|}$

Consider now the subgroups of order four.

Subgroups $R_{(a),(b)}$, $R_{(b),(c)}$, $R_{(b),(ac)}$ cannot have invariant classes, because their invariant classes would also be invariant with respect to $R_{(b)}$, which does not have invariant classes.

The subgroup $R_{(a),(c)} = (I, a, c, ac)$ has no invariant classes. The equation (7) has no solution, indeed the supersymmetric front would satisfy both (8) and (11), which are incompatible.

The other subgroups of RO of order 4 have invariant classes. The solutions of eq. (7) are shown in the table, i.e. the supersymmetric fronts satisfy respectively:

the front $\gamma_{a,bc}$ satisfies:

$$\begin{aligned} -x(t) &= x(t + \pi) \\ y(t) &= y(t + \pi) \\ -\phi(t) &= \phi(t + \pi) \end{aligned} \quad (14)$$

and

$$\begin{aligned} -x(-t) &= x(t) \\ -y(-t) &= y(t) \\ \phi(-t) &= \phi(t) \end{aligned} \quad (15)$$




Fig.16

A supersymmetric front $\gamma_{a,bc}$ with $i = 0$ and $\mu = 2$.




Fig.17

The 2 $R_{(a),(bc)}$ -invariant classes in $\mathcal{N}_{0,|\mu|}$

the front $\gamma_{c,ab}$ satisfies:

$$\begin{aligned}
 -x(t) &= (\cos\alpha)x(t + \pi + \alpha) - (\sin\alpha)y(t + \pi + \alpha) \\
 -y(t) &= (\cos\alpha)y(t + \pi + \alpha) + (\sin\alpha)x(t + \pi + \alpha) \\
 \phi(t) &= \phi(t + \pi + \alpha) + \alpha
 \end{aligned}
 \tag{16}$$

where $\alpha = (2n+1)\frac{\pi}{p}$ (symmetry of order p even), or $\alpha = \frac{2n\pi}{p}$ (symmetry of order p odd).

and

$$\begin{aligned}
 -x(-t) &= x(t) \\
 y(-t) &= y(t) \\
 -\phi(-t) &= \phi(t)
 \end{aligned}
 \tag{17}$$

In fig. 18 examples of the case of no rotation symmetry ($p = 1$) and of rotation symmetry ($p = 2$) are given. An example of the case $p = 3$ is shown in fig. 22.




Fig.18

Supersymmetric fronts $\gamma_{c,ab}$ with $\mu = 0, i = 0$ and $\mu = 0, i = 1$



$i = 1 \quad \mu = 0$	$i = -1 \quad \mu = 0$
 <p data-bbox="226 1263 623 1299">$[\gamma_{c,ab}] = [c\gamma_{c,ab}] = [ab\gamma_{c,ab}] = [abc\gamma_{c,ab}]$</p>	 <p data-bbox="642 1263 1026 1299">$[a\gamma_{c,ab}] = [ac\gamma_{c,ab}] = [b\gamma_{c,ab}] = [bc\gamma_{c,ab}]$</p>

Fig.19

The 2 $R_{(c),(ab)}$ -invariant classes in $\mathcal{N}_{|i|,0}$

the front $\gamma_{ab,bc}$ satisfies:

$$\begin{aligned} -x(-t) &= x(t + \pi) \\ y(-t) &= y(t + \pi) \\ -\phi(-t) &= \phi(t + \pi) \end{aligned} \tag{18}$$

and

$$\begin{aligned} -x(-t) &= x(t) \\ -y(-t) &= y(t) \\ \phi(-t) &= \phi(t) \end{aligned} \tag{19}$$




Fig.20

A supersymmetric front $\gamma_{ab,bc}$ with $i = 0$ and $\mu = 0$




Fig.21

The 2 $R_{(ab),(bc)}$ -invariant classes in $\Omega_{0,0}$

3. ROTATION SYMMETRIES

Definition. A planar front γ is called *p-symmetric* if

$$\tilde{\gamma} = \tau_{\alpha, \beta} \tilde{\gamma}$$

where $\beta = \frac{2\pi}{p}$, $\alpha = \pm\beta$

Proposition. *Every p-symmetric front ($p > 1$) belongs to one of the 4 following symmetry classes:*

$$[\gamma_{ab}], \quad [\gamma_{abc}], \quad [\gamma_c], \quad [\gamma_{c,ab}]$$

Proof. The index i of a p -symmetric front is $i = p \times i_p$ where $2\pi i_p$ is the total rotation angle of the normal vector along a p -th part of the front (for example in the interval $[t, t + \beta]$ of the preimage circle). By hypothesis $\phi(t) = \phi(t + \beta) \pm \beta$ (modulo 2π) so that $2\pi i_p = \phi(t + \beta) - \phi(t)$ cannot be zero.

Thus the symmetries allowed are those with index different from zero. For every type of such symmetries we give examples in fig. 22.

4. OTHER TYPES OF CLASSIFICATION

The above classification contains less refined classifications.

For example, we can consider the classification problem when the front class is taken up to diffeomorphisms not necessarily preserving the orientation of the circle. The answer is thus given by the above table, where the element b is everywhere substituted by the identity. The quotient group (I, a, c, ac) has in this case one invariant class, corresponding of course to the supersymmetric class $[\gamma_{c,ab}] = [\gamma_{ab,bc}]$

Remark. In all such less refined classifications, all subgroups have invariant classes.




Fig.22
 p -symmetric fronts with their RO -symmetries

Example. The involution σ considered in [1,2] acting on the C^1 -immersions of the circle into the plane (the coorientation defined by the normal vector being not considered) has invariant classes.

The classification "up to the change of an orientation" can be also seen as a classification forgetting such orientation. Hence the quotient group (I, a, b, ab) acts also on planar *nonorientable fronts* with semi-integer index (i.e. with an odd number of cusps), which can be defined as projections onto the plane of legendrian immersions of S^1 into the space of non oriented contact elements of the plane $\hat{M}^3 = PT^*\mathbb{R}^2$ (where $M^3 \rightarrow \hat{M}^3$ is the double covering.)

5. DIVERTISSEMENT

Associate to every smooth segment of a oriented and cooriented front in the oriented plane a sign s in the following way: if the pair (orienting tangent vector, coorienting normal vector) defines the positive orientation of the plane (x, y) , then $s = 1$, otherwise $s = -1$.

The sign changes at every cusp of the front.

Define now the length l of a front as the algebraic sum of the lengths of the pieces with their signs. We have associated to the front a real number l (its length).

Remark. The sign of l , if there are cusps, is not an invariant of the front class (see fig.23)




Fig. 23

2 fronts of the same front class and opposite values of the sign of l

However l is invariant in every T -class. The analysis of symmetries, by theorem 2, is made inside T -classes, so that we use the number l as an invariant.

The sign l of a front changes for all the involutions Σ , σ and Θ . This means that the fronts which are invariant with respect to the subgroups $R_{(a)}$, $R_{(b)}$, $R_{(c)}$ and $R_{(abc)}$ have all length zero, and those which are invariant for the other ones can have positive or negative lengths.

We visualize the symmetries of the group RO acting on the fronts using a cube (see fig. 24): the 8 vertices of the cube represent an asymmetric front with index i , Maslov index μ and length l under the action of the 8 elements of RO .




Fig. 24
The cube of the symmetries RO

The cube belongs to 3-dimensional space with coordinates (l, μ, i) so that different values correspond to every vertex.

The planes $l = 0$, $\mu = 0$ and $i = 0$ represent the fronts invariant under the action of the subgroups of order two $R_{(abc)}$, $R_{(ab)}$ and $R_{(bc)}$ respectively (the action is the reflection on these planes).

The rotations by an angle π around the 3 principal axes represent the other 3 subgroups of order two: $R_{(a)}$ ($i = 0, l = 0$), $R_{(c)}$ ($\mu = 0, l = 0$), and $R_{(ac)}$ ($i = 0, \mu = 0$).

The principal axes represent the fronts invariant under the action of the subgroups of order four. Every axis is in fact invariant under the

rotation around itself and under the reflection with respect to the planes containing it.

Remark. In this scheme the subgroups $R_{(b)}$ and $R_{(a,c)}$, which have no invariant classes, leave invariant only the origin.

References

- [1] V.I. Arnold *Invarianty i perestroiki ploskikh frontov*, Trudy Mian, vol. 209, 1995.
- [2] V.I. Arnold *Plane Curves, Their Invariants, Perestroikas and Classifications*, Adv. in Sov. Math., A.M.S. 21, Providence, 1994.

International Centre for Theoretical Physics
Strada Costiera, 11
I-34014 Trieste
ITALIA

Recibido: 4 de Agosto de 1994