

# A capacity approach to the Poincaré inequality and Sobolev imbeddings in variable exponent Sobolev spaces

Petteri HARJULEHTO and Peter HÄSTÖ

Department of Mathematics  
P.O. Box 4 (Yliopistonkatu 5)  
FIN-00014 University of Helsinki, Finland  
petteri.harjulehto@helsinki.fi,  
peter.hasto@helsinki.fi

Recibido: 7 de Mayo de 2003

Aceptado: 10 de Septiembre de 2003

## ABSTRACT

We study the Poincaré inequality in Sobolev spaces with variable exponent. Under a rather mild and sharp condition on the exponent  $p$  we show that the inequality holds. This condition is satisfied e. g. if the exponent  $p$  is continuous in the closure of a convex domain. We also give an essentially sharp condition for the exponent  $p$  as to when there exists an imbedding from the Sobolev space to the space of bounded functions.

*Key words:* Sobolev spaces, variable exponent, Poincaré inequality, Sobolev imbedding, continuity

*2000 Mathematics Subject Classification:* 46E35

## 1. Introduction

There has recently been a surge of interest in Sobolev spaces with variable exponent, cf. [4–7, 9–11, 17, 22]. These spaces, introduced in [17], are the natural generalization of Sobolev spaces to the non-homogeneous situation; they have been used e. g. in modeling electrorheological fluids, see the book of M. Růžička, [22]. Lebesgue and Sobolev spaces with variable exponent share many properties with their classical equivalents, but there is also some crucial differences. For instance the Hardy-Littlewood maximal

---

The second author was supported financially by the Academy of Finland.

operator is bounded on  $L^{p(\cdot)}$  if the exponent is 0-Hölder continuous (i. e. satisfies (10)) and  $1 < \text{ess inf } p \leq \text{ess sup } p < \infty$ , [5]. If the exponent is not 0-Hölder continuous, then the maximal operator need not be bounded on  $L^{p(\cdot)}$ , [21].

The Poincaré inequality, although of great importance in classical non-linear potential theory (especially in metric spaces) has not been previously studied in the case of variable exponent Sobolev spaces. Our first result, Theorem 2.2, is the following: If  $D \subset \mathbb{R}^n$  is smooth domain, say a John domain, and the essential supremum of  $p$  is less than the Sobolev conjugate of the essential infimum of  $p$  then the Poincaré inequality

$$\|u - u_B\|_{L^{p(\cdot)}(D)} \leq C \|\nabla u\|_{L^{p(\cdot)}(D)}$$

holds for every  $u \in W^{1,p(\cdot)}(D)$ , where  $u_B = \int u(x)dx$ . Here the constant  $C$  depends on  $n, p, \text{diam}(D)$  and the John constant of  $D$ . We give an example which shows that the condition for  $p$  is sharp even in a ball. It follows from this that if  $p$  is continuous in the closure of a convex domain then the Poincaré inequality holds (Corollary 2.7).

In classical theory the constant of the Poincaré inequality is  $C \text{diam}(D)$ . It is possible to achieve this also for variable exponent Sobolev spaces, as we prove in Corollary 2.10. The price we have to pay is that the exponent  $p$  has to be 0-Hölder continuous.

Sobolev imbeddings in variable exponent Sobolev spaces have been studied by many authors in the case when  $p$  is less than the dimension, see [6, 9–11]. We give two results in the case when  $p$  is greater than the dimension. We prove a result for continuity of the Sobolev functions, namely that every Sobolev function is continuous if the exponent is locally bounded away from the dimension. We show that if a domain satisfies a uniform interior cone condition and  $p(x) \geq n + f(d(x, \partial G))$  for every  $x$  and a certain increasing function  $f$  then there exists an imbedding from the variable exponent Sobolev space to  $L^\infty$ . Our condition is essentially sharp.

**Notation**

We denote by  $\mathbb{R}^n$  the Euclidean space of dimension  $n \geq 2$ . For  $x \in \mathbb{R}^n$  and  $r > 0$  we denote an open ball with center  $x$  and radius  $r$  by  $B(x, r)$ .

Let  $A \subset \mathbb{R}^n$  and  $p: A \rightarrow [1, \infty)$  be a measurable function (called a *variable exponent* on  $A$ ). We define  $p_A^+ = \text{ess sup}_{x \in A} p(x)$  and  $p_A^- = \text{ess inf}_{x \in A} p(x)$ . If  $A = \mathbb{R}^n$  we write  $p^+ = p_{\mathbb{R}^n}^+$  and  $p^- = p_{\mathbb{R}^n}^-$ .

Let  $\Omega \subset \mathbb{R}^n$  be an open set. We define the *generalized Lebesgue space*  $L^{p(\cdot)}(\Omega)$  to consist of all measurable functions  $u: \Omega \rightarrow \mathbb{R}$  such that

$$\varrho_{p(\cdot)}(\lambda u) = \int_{\Omega} |\lambda u(x)|^{p(x)} dx < \infty$$

for some  $\lambda > 0$ . The function  $\varrho_{p(\cdot)}: L^{p(\cdot)}(\Omega) \rightarrow [0, \infty)$  is called the *modular* of the space  $L^{p(\cdot)}(\Omega)$ . One can define a norm, the so-called *Luxemburg norm*, on this space by the formula  $\|u\|_{p(\cdot)} = \inf\{\lambda > 0 : \varrho_{p(\cdot)}(u/\lambda) \leq 1\}$ . Notice that if  $p \equiv p_0$  then

$L^{p(\cdot)}(\Omega)$  is the classical Lebesgue space, so there is no danger of confusion with the new notation.

The *generalized Sobolev space*  $W^{1,p(\cdot)}(\Omega)$  is the space of measurable functions  $u: \Omega \rightarrow \mathbb{R}$  such that  $u$  and the absolute value of the distributional gradient  $\nabla u = (\partial_1 u, \dots, \partial_n u)$  are in  $L^{p(\cdot)}(\Omega)$ . The function  $\varrho_{1,p(\cdot)}: W^{1,p(\cdot)}(\Omega) \rightarrow [0, \infty)$  is defined as  $\varrho_{1,p(\cdot)}(u) = \varrho_{p(\cdot)}(u) + \varrho_{p(\cdot)}(|\nabla u|)$ . The norm  $\|u\|_{1,p(\cdot)} = \|u\|_{p(\cdot)} + \|\nabla u\|_{p(\cdot)}$  makes  $W^{1,p(\cdot)}(\mathbb{R}^n)$  a Banach space.

See [17] for basic properties of variable exponent Lebesgue and Sobolev spaces.

## 2. The Poincaré inequality

In this section we give a relatively mild condition on the exponent for the Poincaré inequality to hold. We also show that this condition is, in a certain sense, the best possible. For Sobolev functions with zero boundary values the Poincaré inequality was given in [10, Lemma 3.1] and considerably generalized in [14].

Recall the following well known Sobolev-Poincaré inequality. By  $q^*$  we denote the Sobolev conjugate of  $q < n$ ,  $q^* = nq/(n - q)$ .

**Lemma 2.1.** *Let  $D \subset \mathbb{R}^n$  be a bounded John domain. Let  $1 \leq p < n$  and  $p \leq q \leq p^*$  be fixed exponents. Then*

$$\|u - u_D\|_q \leq C(n, p, \lambda) |D|^{1/n+1/q-1/p} \|\nabla u\|_p$$

for all functions  $u \in W^{1,p}(D)$ , where  $\lambda$  is the John constant.

If  $p \geq n$  and  $q < \infty$  then

$$\|u - u_D\|_q \leq C(n, q, \lambda) |D|^{1/n+1/q-1/p} \|\nabla u\|_p$$

for all functions  $u \in W^{1,p}(D)$ .

*Proof.* The case  $p < n$  and  $q = p^*$  is by B. Bojarski [3, (6.6)]. The case  $q < p^*$  follows from this by standard arguments: we choose  $s \in [1, n)$  such that  $s^* = q$  (or  $s = 1$  if  $q < 1^*$ ). By Hölder's inequality and Bojarski's result we obtain

$$\begin{aligned} \left( \int_D |u - u_D|^q dx \right)^{\frac{1}{q}} &\leq |D|^{-\frac{1}{s^*}} \left( \int_D |u - u_D|^{s^*} dx \right)^{\frac{1}{s^*}} \leq C |D|^{-\frac{1}{s^*}} \left( \int_D |\nabla u|^s dx \right)^{\frac{1}{s}} \\ &= C |D|^{\frac{1}{s} - \frac{1}{s^*}} \left( \int_D |\nabla u|^s dx \right)^{\frac{1}{s}} \leq C |D| \left( \int_D |\nabla u|^p dx \right)^{\frac{1}{p}}, \end{aligned}$$

which is clearly equivalent to the inequalities in the theorem. □

**Theorem 2.2.** *Let  $D \subset \mathbb{R}^n$  be a bounded John domain, with constant  $\lambda$ . If  $p_D^+ \leq (p_D^-)^*$  or  $p_D^- \geq n$  and  $p_D^+ < \infty$  then there exists a constant  $C = C(n, p_D^-, p_D^+, \lambda)$  such that*

$$\|u - u_D\|_{p(\cdot)} \leq C(1 + |D|)^2 |D|^{\frac{1}{n} + \frac{1}{p_D^+} - \frac{1}{p_D^-}} \|\nabla u\|_{p(\cdot)} \tag{1}$$

for every  $u \in W^{1,p(\cdot)}(D)$ .

*Proof.* Assume first that  $p_D^+ \leq (p_D^-)^*$ . Since  $p(x) \leq p_D^+ \leq (p_D^-)^*$  we obtain by [17, Theorem 2.8] and Lemma 2.1 that

$$\begin{aligned} \|u - u_D\|_{p(\cdot)} &\leq (1 + |D|) \|u - u_D\|_{p_D^+} \\ &\leq C(n, p_D^-, \lambda) (1 + |D|) |D|^{\frac{1}{n} + \frac{1}{p_D^+} - \frac{1}{p_D^-}} \|\nabla u\|_{p_D^-} \\ &\leq C(n, p_D^-, \lambda) (1 + |D|)^2 |D|^{\frac{1}{n} + \frac{1}{p_D^+} - \frac{1}{p_D^-}} \|\nabla u\|_{p(\cdot)}. \end{aligned}$$

The case  $p_D^- \geq n$  is similar, the only difference is that the constant in the second inequality in the above chain of inequalities is  $C(n, p_D^+, \lambda)$ . □

*Remark 2.3.* John domains are almost the right class of irregular domains for the classical Sobolev-Poincaré inequality, see [3], [1] and [2, Theorem 4.1].

Previous results on Sobolev imbeddings in the variable exponent setting have been derived in domains whose boundary is locally a graph of a Lipschitz continuous function, see [9–11]. It is therefore of interest to note that every domain, whose boundary is locally the graph of a Lipschitz continuous function, is a John domain, see [19]. In particular every ball is a John domain.

If  $D$  is a ball in Theorem 2.2, then the constant in inequality (1) is the classical Sobolev-Poincaré inequality in a ball, see for example [18, Corollary 1.64, p. 38].

The next example shows that if  $p_D^- < n$  and  $p_D^+ > (p_D^-)^*$  then there need not exist a constant  $C > 0$  such that inequality (1) holds for every  $u \in W^{1,p(\cdot)}(D)$ .

Recall that the *variational capacity* for fixed  $p$ ,  $\text{cap}_p(E, F; D)$ , is defined for sets  $E, F$  and open  $D$  by

$$\text{cap}_p(E, F; D) = \inf_{u \in L(E, F; D)} \int_D |\nabla u|^p dx,$$

where  $L(E, F; D)$  is the set of continuous functions  $u$  that satisfy  $u|_{E \cap D} = 1$ ,  $u|_{F \cap D} = 0$  and  $|\nabla u| \in L^{p(\cdot)}(D)$ . We use the short-hand notation  $\text{cap}(E, F)$  for  $\text{cap}(E, F; \mathbb{R}^n)$ , similarly for  $L(E, F)$ . For more information on capacities see [15, Chapter 2] or [20]. The following lemma will be used several times to estimate the gradient of variable exponent functions.

**Lemma 2.4** ([15, Example 2.12, p. 35]). *For fixed  $p \neq 1, n$ , arbitrary  $x \in \mathbb{R}^n$  and  $R > r > 0$  we have*

$$\text{cap}_p(\mathbb{R}^n \setminus B(x, R), B(x, r)) = \omega_{n-1} \left| \frac{p-n}{p-1} \right|^{p-1} \left| R^{(p-n)/(p-1)} - r^{(p-n)/(p-1)} \right|^{1-p}.$$

**Example 2.5.** Our aim is construct a sequence of functions in  $B = B(0, 1) \subset \mathbb{R}^2$  for which the constant in the Poincaré inequality (1) goes to infinity. Let  $B_i = B(2^{-i}e_1, \frac{1}{4}2^{-i}) \subset \mathbb{R}^2$  and  $B'_i = B(2^{-i}e_1, \frac{1}{8}2^{-i^2}) \subset \mathbb{R}^2$  for every  $i = 1, 2, \dots$  and let  $1 < p_1 < 2$ . Choose a function  $u_i \in C_0^\infty(B_i)$  with  $u_i|_{B'_i} = 1$  be such that

$$\left( 2 \text{cap}_{p_1}(B'_i, \mathbb{R}^2 \setminus B_i) \right)^{\frac{1}{p_1}} \geq \|\nabla u_i\|_{L^{p_1}(B_i)}. \tag{2}$$

Let  $p_2 > 2$  and define  $p(x) = p_1\chi_{B_i \setminus B'_i}(x) + p_2\chi_{B'_i}(x)$  for  $x \in B$  with positive first coordinate. Since  $\nabla u_i = 0$  in  $B'_i$  we obtain

$$\|\nabla u_i\|_{L^{p(\cdot)}(B_i)} = \|\nabla u_i\|_{L^{p_1}(B_i)}. \tag{3}$$

Let  $\tilde{B}_i = B(-2^{-i}e_1, \frac{1}{4}2^{-i})$ . We extend  $u_i$  to  $B$  as an odd function of the first coordinate in  $\tilde{B}_i$  and by zero elsewhere. We also extend  $p$  to  $B$  as an even function of the first coordinate. We denote the extensions by  $\tilde{u}_i$  and  $\tilde{p}$ . By (2) and (3) we obtain

$$2^{1+\frac{1}{p_1}} \left( \text{cap}_{p_1}(B'_i, \mathbb{R}^2 \setminus B_i) \right)^{\frac{1}{p_1}} \geq \|\nabla \tilde{u}_i\|_{L^{\tilde{p}(\cdot)}(B)}.$$

By Lemma 2.4 this yields

$$\|\nabla \tilde{u}_i\|_{L^{\tilde{p}(\cdot)}(B)} \leq C(p_1) \left| \frac{1}{4}2^{-i\frac{p_1-2}{p_1-1}} - \frac{1}{8}2^{-i^2\frac{p_1-2}{p_1-1}} \right|^{\frac{1-p_1}{p_1}}. \tag{4}$$

For large  $i$  the right hand side is approximately equal to  $C(p_1)2^{-i^2\frac{2-p_1}{p_1}}$ .

Since  $(\tilde{u}_i)_B = 0$ , we obtain

$$\|\tilde{u} - (\tilde{u}_i)_B\|_{L^{\tilde{p}(\cdot)}(B)} = \|\tilde{u}\|_{L^{\tilde{p}(\cdot)}(B)} \geq |B'_i|^{\frac{1}{p_2}} \approx 2^{-i^2\frac{2}{p_2}}. \tag{5}$$

By inequalities (4) and (5) we find that

$$\frac{\|\tilde{u} - (\tilde{u}_i)_B\|_{L^{\tilde{p}(\cdot)}(B)}}{\|\nabla \tilde{u}_i\|_{L^{\tilde{p}(\cdot)}(B)}} \geq C(p_1)2^{i^2(\frac{2}{p_1}-1-\frac{2}{p_2})} \rightarrow \infty$$

as  $i \rightarrow \infty$  if  $\frac{2}{p_1} - 1 - \frac{2}{p_2} > 0$ , that is, if  $p_2 > \frac{2p_1}{2-p_1} = p_1^*$ .

We next show that the condition  $p_D^\dagger \leq (p_D^-)^*$  in Theorem 2.2 can be replaced by a set of local conditions.

**Theorem 2.6.** *Let  $D \subset \mathbb{R}^n$  be a bounded John domain. Assume that there exist John domains  $G_i$ ,  $i = 1, \dots, j$ , so that  $G_i \subset D$  for every  $i$ ,  $D = \cup_{i=1}^j G_i$  and either  $p_{G_i}^+ \leq (p_{G_i}^-)^*$  or  $p_{G_i}^- \geq n$  for every  $i$ . Then there exists a constant  $C > 0$  such that*

$$\|u - u_D\|_{p(\cdot)} \leq C \|\nabla u\|_{p(\cdot)} \tag{6}$$

for every  $u \in W^{1,p(\cdot)}(D)$ . The constant  $C$  depends on  $n$ ,  $\text{diam}(D)$ ,  $|G_i|$ ,  $p$  and the John constants of  $D$  and  $G_i$ ,  $i = 1, \dots, j$ .

*Proof.* Using the triangle inequality of the norm we obtain

$$\begin{aligned} \|u - u_D\|_{L^{p(\cdot)}(D)} &\leq \sum_{i=1}^j \|u - u_D\|_{L^{p(\cdot)}(G_i)} \\ &\leq \sum_{i=1}^j \|u - u_{G_i}\|_{L^{p(\cdot)}(G_i)} + \sum_{i=1}^j \|u_D - u_{G_i}\|_{L^{p(\cdot)}(G_i)}. \end{aligned} \tag{7}$$

We estimate the first part of the sum using Theorem 2.2. This yields

$$\begin{aligned} \|u - u_{G_i}\|_{L^{p(\cdot)}(G_i)} &\leq C(n, p_{G_i}, |G_i|, \lambda_i) \|\nabla u\|_{L^{p(\cdot)}(G_i)} \\ &\leq C(n, p_{G_i}, |G_i|, \lambda_i) \|\nabla u\|_{L^{p(\cdot)}(D)} \end{aligned} \tag{8}$$

for every  $i = 1, \dots, j$ . Here  $\lambda_i$  is the John constant of  $G_i$ . We next estimate the second part of the sum in (7) using the classical Poincaré inequality for the third inequality. We obtain

$$\begin{aligned} \|u_D - u_{G_i}\|_{L^{p(\cdot)}(G_i)} &\leq \|1\|_{L^{p(\cdot)}(G_i)} \int_{G_i} |u(x) - u_D| dx \\ &\leq \|1\|_{L^{p(\cdot)}(G_i)} |G_i|^{-1} \int_D |u(x) - u_D| dx \\ &\leq C(n, \text{diam}(D), \lambda) |G_i|^{-1} \|1\|_{L^{p(\cdot)}(G_i)} \|\nabla u\|_{L^1(D)} \\ &\leq C(n, \text{diam}(D), \lambda) (1 + |D|) |G_i|^{-1} \|1\|_{L^{p(\cdot)}(G_i)} \|\nabla u\|_{L^{p(\cdot)}(D)} \end{aligned} \tag{9}$$

for every  $i = 1, \dots, j$ . Here  $\lambda$  is the John constant of  $D$ . Now inequality (6) follows by inequalities (7), (8) and (9). □

**Corollary 2.7.** *Let  $D \subset \mathbb{R}^n$  be a bounded convex domain and let  $p: \overline{D} \rightarrow [1, \infty)$  be a continuous exponent. Then there exists a constant  $C > 0$  such that*

$$\|u - u_D\|_{p(\cdot)} \leq C \|\nabla u\|_{p(\cdot)}$$

for every  $u \in W^{1,p(\cdot)}(\mathbb{R}^n)$ .

*Proof.* Since  $p$  is continuous we find for every  $x \in \bar{D}$  a constant  $r(x) > 0$  such that either

$$p_{B(x,r(x)) \cap D}^+ \leq (p_{B(x,r(x)) \cap D}^-)^* \quad \text{or} \quad p_{B(x,r(x)) \cap D}^- \geq n.$$

Since  $\bar{D}$  is compact it is possible to find finite covering of  $D$  with balls  $B(x, r(x))$ . It is easy to see that each  $B(x, r(x)) \cap D$  is a John domain and hence the corollary follows by Theorem 2.6.  $\square$

Sometimes it is useful to have better control over the constant in the Poincaré inequality as the domain  $D$  changes than we have in (1). In the fixed exponent case the constant of the Poincaré inequality is  $C \text{diam}(D)$ . We show that this kind of constant is also possible for variable exponent Sobolev spaces. The price we have to pay for this is that the exponent  $p$  has to satisfy a much stronger condition in Theorem 2.8 than in Theorem 2.2; in Theorem 2.2 the exponent  $p$  could be discontinuous even in every point, but in Theorem 2.8 the exponent is 0-Hölder continuous.

**Theorem 2.8.** *Let  $D \subset \mathbb{R}^n$  be a bounded uniform domain. Let  $p: D \rightarrow \mathbb{R}$  be such that  $1 < p_D^- \leq p_D^+ < \infty$ . Assume that there exists a constant  $C > 0$  such that*

$$|p(x) - p(y)| \leq \frac{C}{-\log|x - y|} \tag{10}$$

for every  $x, y \in D$  with  $|x - y| \leq \frac{1}{2}$ . Then the inequality

$$\|u - u_D\|_{p(\cdot)} \leq C \text{diam}(D) \left( 1 + \max \left\{ |D|^{1/p_D^+ - 1/p_D^-}, |D|^{1/p_D^- - 1/p_D^+} \right\} \right) \|\nabla u\|_{p(\cdot)}, \tag{11}$$

holds for every  $u \in W^{1,p(\cdot)}(D)$ . Here the constant  $C$  depends on the dimension  $n$ , the uniform constant of  $D$  and  $p$ .

*Proof.* Since  $W_0^{1,p(\cdot)}(D) \hookrightarrow W^{1,1}(D)$  we obtain as in the proof of [12, Theorem 11] for every  $u \in W^{1,p(\cdot)}(D)$  that

$$|u(x) - u(y)| \leq C|x - y|(\mathbb{M}\nabla u(x) + \mathbb{M}\nabla u(y)) \tag{12}$$

for almost every  $x, y \in D$ . Here  $\mathbb{M}$  is the Hardy-Littlewood maximal operator:

$$\mathbb{M}\nabla u(x) = \sup_{r>0} \int_{B(x,r)} |\nabla u(y)| dy,$$

with the understanding that  $\nabla u = 0$  outside  $D$ . The constant  $C$  depends on the dimension  $n$  and the uniform constants of  $D$ .

Integrating inequality (12) over  $y$  we obtain

$$\begin{aligned} \left| u(x) - \int_D u(y) dy \right| &\leq \int_D |u(x) - u(y)| dy \\ &\leq C \text{diam}(D) \left( \mathbb{M}\nabla u(x) + \int_D \mathbb{M}\nabla u(y) dy \right). \end{aligned}$$

By Hölder's inequality [17, Theorem 2.1] this yields

$$|u(x) - u_D| \leq C \operatorname{diam}(D) \left( \mathbb{M}\nabla u(x) + \frac{C(p)\|1\|_{L^{p'(\cdot)}(D)}}{|D|} \|\mathbb{M}\nabla u\|_{p(\cdot)} \right).$$

Since the previous inequality holds point-wise, it is clear that we have an inequality also for the Lebesgue norms of both sides:

$$\begin{aligned} \|u - u_D\|_{p(\cdot)} &\leq C \operatorname{diam}(D) \left( \|\mathbb{M}\nabla u\|_{p(\cdot)} + \frac{C}{|D|} \|1\|_{p'(\cdot)} \|1\|_{p(\cdot)} \|\mathbb{M}\nabla u\|_{p(\cdot)} \right) \\ &\leq C \operatorname{diam}(D) \left( 1 + |D|^{-1} \max\{|D|^{1+1/p_D^+ - 1/p_D^-}, |D|^{1+1/p_D^- - 1/p_D^+}\} \right) \|\mathbb{M}\nabla u\|_{p(\cdot)} \end{aligned}$$

By [5, Theorem 3.5] (see also [7, Remark 2.2]) the Hardy-Littlewood maximal operator is bounded, and so we obtain

$$\|u - u_D\|_{p(\cdot)} \leq C \operatorname{diam}(D) \left( 1 + \max\left\{|D|^{1/p_D^+ - 1/p_D^-}, |D|^{1/p_D^- - 1/p_D^+}\right\} \right) \|\nabla u\|_{p(\cdot)},$$

where the constant  $C$  depends on the dimension  $n$ , the uniform constant of  $D$  and  $p$ . □

*Remark 2.9.* We refer to [19] for basic properties of uniform domains: Every uniform domain is a John domain. Every domain, whose boundary is locally a graph of a Lipschitz continuous function, is a uniform domain. In particular if  $D$  is a ball then the constant in (11) depends on the dimension  $n$  and  $p$ .

**Corollary 2.10.** *Let  $p$  be as in the previous theorem. If  $B$  is a ball with  $|B| \leq 1$  then*

$$\|u - u_B\|_{p(\cdot)} \leq C \operatorname{diam}(B) \|\nabla u\|_{p(\cdot)},$$

where the constant  $C$  does not depend on  $B$ .

*Proof.* Since  $|B| \leq 1$  we have

$$\max\left\{|B|^{1/p_B^+ - 1/p_B^-}, |B|^{1/p_B^- - 1/p_B^+}\right\} = |B|^{1/p_B^+ - 1/p_B^-}.$$

Since  $p$  is 0-Hölder continuous, (10), we obtain by [5, Lemma 3.2] that there exists a constant  $C > 0$ , depending only on the dimension  $n$  and the constant in (10), such that  $|B|^{1/p_B^+ - 1/p_B^-} \leq C$  for every ball  $B$ . Hence  $|B| \leq 1$  implies that the constant in (11) is less than  $C \operatorname{diam}(B)$ . □



### 3. Continuity

The functions in the classical Sobolev space  $W^{1,p}$  are continuous if  $p > n$ . In this section we consider when functions in variable exponent Sobolev space are continuous.

**Theorem 3.1.** *Suppose that  $p > n$  is locally bounded away from  $n$  in  $D$ . Then  $W^{1,p(\cdot)}(D) \subset C(D)$ .*

*Proof.* Let  $x \in D$  and consider the ball  $B = B(x, \delta(x)/2)$ . Define  $q = \text{ess inf}_{y \in B} p(y)$ . Then, by [17, Theorem 2.8],

$$W^{1,p(\cdot)}(B) \hookrightarrow W^{1,q}(B) \subset C(B).$$

Therefore every function in  $W^{1,p(\cdot)}(D)$  is continuous at  $x$ , and since  $x$  was arbitrary, the claim follows.  $\square$

The following corollary is immediate.

**Corollary 3.2.** *Suppose that  $p$  is continuous in  $D$ . Then  $W^{1,p(\cdot)}(D) \subset C(D)$  if  $p(x) > n$  for every  $x \in D$ .*

We next use a classical example to show that the assumption that  $p$  is locally bounded away from  $n$  in  $D$  is not superfluous when  $p$  is not continuous.

**Example 3.3.** Let  $B = B(0, 1/16)$ ,  $\varepsilon > 0$  and suppose that

$$p(x) \leq \bar{p}(|x|) = n + (n - 1 - \varepsilon) \frac{\log_2 \log_2(1/|x|)}{\log_2(1/|x|)}$$

for  $x \in B \setminus \{0\}$  and  $p(0) > n$ . We show that then  $W^{1,p(\cdot)}(B) \not\subset C(B)$ .

Define  $u(x) = \cos(\log_2 |\log_2 |x||)$  for  $x \in B \setminus \{0\}$  and  $u(0) = 0$ . Clearly  $u$  is not continuous at the origin. So we have to show that  $u \in W^{1,p(\cdot)}(B)$ . It is clear that  $u$  has partial derivatives, except at the origin.

Since  $u$  is bounded it follows that  $u \in L^{p(\cdot)}(B)$ . We next estimate the gradient:

$$|\nabla u(x)| = \left| \sin(\log_2 |\log_2 |x||) \cdot \frac{1}{|x| \log_2 |x|} \right| \leq \left| \frac{1}{|x| \log_2 |x|} \right|.$$

We therefore find that

$$\begin{aligned} \int_B |\nabla u(x)|^{p(x)} dx &\leq \int_B \frac{dx}{(|x| |\log_2 |x||)^{p(x)}} \\ &= \omega_{n-1} \int_0^{1/16} \frac{r^{n-1} dr}{(r |\log_2 r|)^{\bar{p}(r)}} \\ &= \omega_{n-1} \sum_{i=5}^{\infty} \int_{2^{-i-1}}^{2^{-i}} \frac{r^{n-1} dr}{(r |\log_2 r|)^{\bar{p}(r)}}. \end{aligned}$$

Since  $1/(r|\log_2 r|) > 1$  we may increase the exponent  $\bar{p}$  for an upper bound. In the annulus  $B(0, 2^{-i}) \setminus B(0, 2^{-i-1})$  we have  $i \leq \log_2(1/|x|) \leq i + 1$ . Since  $y \rightarrow \log_2(y)/y$  is decreasing we find that

$$\bar{p}(x) \leq n + (n - 1 - \varepsilon) \frac{\log_2 i}{i}$$

in the same annulus. We can therefore continue our previous estimate by

$$\begin{aligned} \int_B |\nabla u(x)|^{p(x)} dx &\leq \sum_{i=5}^{\infty} \int_{2^{-i-1}}^{2^{-i}} \frac{r^{n-1} dr}{(r|\log_2 r|)^{n+(n-1-\varepsilon)\log_2(i)/i}} \\ &\leq C \sum_{i=5}^{\infty} \int_{2^{-i-1}}^{2^{-i}} \frac{2^{-i(n-1)} dr}{(i2^{-i})^{n+(n-1-\varepsilon)\log_2(i)/i}} \\ &= C \sum_{i=5}^{\infty} 2^{(n-1-\varepsilon)\log_2(i)} i^{-n-(n-1-\varepsilon)\log_2(i)/i} \\ &= C \sum_{i=5}^{\infty} i^{-1-\varepsilon} i^{-(n-1-\varepsilon)\log_2(i)/i} \leq C \sum_{i=5}^{\infty} i^{-1-\varepsilon} < \infty. \end{aligned}$$

### 4. Sobolev imbedding theorems

We start by introducing a relative variational  $p(\cdot)$ -pseudocapacity, and proving some basic properties for it. This capacity is quite similar to the Sobolev  $p(\cdot)$ -capacity studied by P. Harjulehto, P. Hästö, M. Koskenoja and S. Varonen in [13].

Let  $F, E \subset \mathbb{R}^n$  be closed disjoint sets and  $D$  be a domain in  $\mathbb{R}^n$ . The *variational  $p(\cdot)$ -pseudocapacity* is defined as

$$\psi_{p(\cdot)}(F, E; D) = \inf_{u \in L(F, E; D)} \|\nabla u\|_{L^{p(\cdot)}(D)},$$

where  $L(F, E; D)$  is as before (see Section 2). For  $L(F, E; D) = \emptyset$  we define  $\psi_{p(\cdot)}(F, E; D) = \infty$ . We write  $L(E, x; D)$  for  $L(F, \{x\}; D)$  etc.

*Remark 4.1.* Including  $C(D)$  in the definition of the capacity is somewhat strange in this context, since we do not, in general, know whether continuous functions are dense in  $W^{1,p(\cdot)}(D)$ , but see [8]. However, since we are interested in the case when  $p > n$ , the assumption makes sense, by Theorem 3.1.

The reason for calling the function  $\psi_{p(\cdot)}(F, E; D)$  a pseudocapacity is that it is defined as a capacity but using the norm instead of the modular. This corresponds to introducing an exponent  $1/p$  to the capacity in the fixed exponent case. Because of this we cannot expect the pseudocapacity to have all the usual properties of a capacity. It nevertheless has many of them:

**Theorem 4.2.** *Let  $F, E \subset \mathbb{R}^n$  be closed sets and  $D$  be a domain in  $\mathbb{R}^n$ . Then the set function  $(F, E) \mapsto \psi_{p(\cdot)}(F, E; D)$  has the following properties:*

- (i)  $\psi_{p(\cdot)}(\emptyset, E; D) = 0$ .
- (ii)  $\psi_{p(\cdot)}(F, E; D) = \psi_{p(\cdot)}(E, F; D)$ .
- (iii) *Outer regularity, i. e.  $\psi_{p(\cdot)}(F, E_1; D) \leq \psi_{p(\cdot)}(F, E_2; D)$ .*
- (iv) *If  $E$  is a subset of  $\mathbb{R}^n$ , then*

$$\psi_{p(\cdot)}(F, E; D) = \inf_{\substack{E \subset U \\ U \text{ open}}} \psi_{p(\cdot)}(F, U; D).$$

- (v) *If  $K_1 \supset K_2 \supset \dots$  are compact, then*

$$\lim_{i \rightarrow \infty} \psi_{p(\cdot)}(F, K_i; D) = \psi_{p(\cdot)}\left(F, \bigcap_{i=1}^{\infty} K_i; D\right).$$

- (vi) *Suppose that  $p > n$  is locally bounded away from  $n$ . If  $E_i \subset \mathbb{R}^n$  for every  $i = 1, 2, \dots$ , then*

$$\psi_{p(\cdot)}\left(F, \bigcup_{i=1}^{\infty} E_i; D\right) \leq \sum_{i=1}^{\infty} \psi_{p(\cdot)}(F, E_i; D).$$

*Proof.* Assertion (i) is clear since we may use a constant function. Assertion (ii) is clear since if  $u \in L(F, E; D)$  then  $1 - u \in L(E, F; D)$ . Assertion (iii) follows since  $L(F, E_2; D) \subset L(F, E_1; D)$ .

Next we prove (iv). It is clear that

$$\psi_{p(\cdot)}(F, E; D) \leq \inf_{\substack{E \subset U \\ U \text{ open}}} \psi_{p(\cdot)}(F, U; D).$$

Let  $\varepsilon > 0$ . Assume that  $u \in L(F, E; D)$  is such that

$$\|\nabla u\|_{p(\cdot)} \leq \psi_{p(\cdot)}(F, E; D) + \varepsilon.$$

Since  $u$  is continuous,  $\{u > 1 - \varepsilon\}$  is an open set containing  $E$ . Hence we obtain

$$\begin{aligned} \inf_{\substack{E \subset U \\ U \text{ open}}} \psi_{p(\cdot)}(F, U; D) &\leq \psi_{p(\cdot)}(F, \{u > 1 - \varepsilon\}; D) \\ &\leq \left\| \nabla \min\left\{\frac{u}{1 - \varepsilon}, 1\right\} \right\|_{p(\cdot)} \leq (1 - \varepsilon)^{-1} \|\nabla u\|_{p(\cdot)} \\ &\leq (1 - \varepsilon)^{-1} \psi_{p(\cdot)}(F, E; D) + \frac{\varepsilon}{1 - \varepsilon}. \end{aligned}$$

Letting  $\varepsilon \rightarrow 0$  yields assertion (iv).

We then prove (v). It is clear that

$$\psi_{p(\cdot)}(F, \cap_{i=1}^{\infty} K_i; D) \leq \lim_{i \rightarrow \infty} \psi_{p(\cdot)}(F, K_i; D)$$

Let  $\varepsilon > 0$ . Assume that  $u \in L(F, \cap_{i=1}^{\infty} K_i; D)$  is such that

$$\|\nabla u\|_{p(\cdot)} \leq \psi_{p(\cdot)}(F, \cap_{i=1}^{\infty} K_i; D) + \varepsilon.$$

When  $i$  is large the set  $K_i$  lies in the closed set  $\{u \geq 1 - \varepsilon\}$ ; therefore

$$\begin{aligned} \lim_{i \rightarrow \infty} \psi_{p(\cdot)}(F, K_i; D) &\leq \psi_{p(\cdot)}(F, \{u \geq 1 - \varepsilon\}; D) \\ &\leq \left\| \nabla \min\left\{\frac{u}{1 - \varepsilon}, 1\right\} \right\|_{p(\cdot)} \leq (1 - \varepsilon)^{-1} \|\nabla u\|_{p(\cdot)} \\ &\leq (1 - \varepsilon)^{-1} \psi_{p(\cdot)}(F, \cap_{i=1}^{\infty} K_i; D) + \frac{\varepsilon}{1 - \varepsilon}. \end{aligned}$$

Letting  $\varepsilon \rightarrow 0$  yields assertion (v).

To prove (vi) let  $\varepsilon > 0$  and choose functions  $u_i \in L(F, E_i; D)$  such that

$$\|\nabla u_i\|_{p(\cdot)} \leq \psi_{p(\cdot)}(F, E_i; D) + \varepsilon/2^i,$$

for  $i = 1, \dots$ . Let  $v_i = u_1 + \dots + u_i$ . Then  $(v_i)$  is a Cauchy sequence, and so it converges to a function  $v \in W^{1,p(\cdot)}(D)$ . Define  $\tilde{v}(x) = \min\{v(x), 1\}$ , so that  $|\tilde{v}| \in L^{p(\cdot)}(D)$  by [13, Theorem 2.2]. It is clear that  $\tilde{v}|_{F \cap D} = 0$  and  $\tilde{v}|_{E \cap D} = 1$ , where  $E = \cup E_i$ . Since  $p > n$  is locally bounded away from  $n$ , it follows from Theorem 3.1 that every function in  $W^{1,p(\cdot)}(D)$  is continuous, and so we have  $\tilde{v} \in L(F, \cup E_i; D)$ , from which the claim easily follows, since

$$\|\nabla \tilde{v}\|_{p(\cdot)} \leq \sum_{i=1}^{\infty} \|\nabla u_i\|_{p(\cdot)} \leq \sum_{i=1}^{\infty} \psi_{p(\cdot)}(F, E_i; D) + \varepsilon. \quad \square$$

Using the pseudocapacity we can start our study of Sobolev-type imbeddings. The following result is the direct generalization of [20, 5.1.1, Theorem 1].

**Theorem 4.3.** *If  $p^+ < \infty$ , then the following two conditions are equivalent:*

- (i)  $W^{1,p(\cdot)}(D) \cap C(D) \hookrightarrow L^\infty(D)$ .
- (ii) *There exist  $r, k > 0$  such that  $\psi_{p(\cdot)}(\overline{D} \setminus B(x, r), x; D) \geq k$  for every  $x \in D$ .*

*Proof.* Suppose that (2) holds, with constants  $r, k > 0$ . Let  $u \in W^{1,p(\cdot)}(D) \cap C(D)$  and let  $y \in D$  be a point with  $u(y) \neq 0$ . Fix a function  $\eta \in C_0^\infty(B(0, 1))$  with  $0 \leq \eta \leq 1$  and  $\eta(0) = 1$ . Define  $v(x) = \eta((x - y)/r)u(x)/u(y)$ . It is clear that  $v \in W^{1,p(\cdot)}(D)$

and since  $v(y) = 1$  and  $v(x) = 0$  for  $x \notin B(y, r)$  we see that  $v \in L(\overline{D} \setminus B(y, r), y; D)$ . It follows that

$$k \leq \psi_{p(\cdot)}(\overline{D} \setminus B(y, r), y; D) \leq \|\nabla v\|_{p(\cdot)}.$$

Then we calculate that

$$\begin{aligned} k|u(y)| &\leq \|\nabla(u(x)\eta((x-y)/r))\|_{p(x)} \\ &\leq \sup_{x \in D} \eta(x)\|\nabla u\|_{p(\cdot)} + \frac{1}{r} \sup_{x \in D} \nabla \eta(x)\|u\|_{p(\cdot)} \\ &\leq \max \left\{ \sup_{x \in D} \eta(x), \frac{1}{r} \sup_{x \in D} \nabla \eta(x) \right\} \|u\|_{1,p(\cdot)}, \end{aligned}$$

so that  $|u(y)|$  is bounded by a constant independent of  $y$ .

Suppose conversely that (1) holds and let  $C$  be a constant such that  $\|u\|_\infty \leq C\|u\|_{1,p(\cdot)}$  for all  $u \in W^{1,p(\cdot)}(D)$ . For functions in  $v \in L(\overline{D} \setminus B(x, r), x; D)$  this gives

$$1 = \|v\|_\infty \leq C\|v\|_{1,p(\cdot)} \leq C(\|\chi_{B(x,r)}\|_{p(\cdot)} + \|\nabla v\|_{p(\cdot)}).$$

Since  $p^+ < \infty$  we can choose  $r$  small enough that  $\|\chi_{B(x,r)}\|_{p(\cdot)} \leq 1/(2C)$ . For such  $r$  we have  $\|\nabla v\|_{p(\cdot)} \geq 1/(2C)$ . It follows that

$$\psi_{p(\cdot)}(\overline{D} \setminus B(x, r), x; D) = \inf_{u \in L(\overline{D} \setminus B(x,r), x; D)} \|\nabla u\|_{p(\cdot)} \geq 1/(2C)$$

for the same  $r$ . □

*Remark 4.4.* Since we do not know whether  $C^\infty(D)$  is dense in  $W^{1,p(\cdot)}(D)$  we have only proved the theorem for continuous functions in  $W^{1,p(\cdot)}(D)$ . If  $p$  is such that  $C(D)$  is dense in  $W^{1,p(\cdot)}(D)$ , for instance if  $p$  is locally bounded above  $n$ , then we may replace condition (1) by  $W^{1,p(\cdot)}(D) \hookrightarrow L^\infty(D)$ .

Define  $D = B(1/16) \setminus \{0\}$  and let  $p$  be as in Example 3.3. Then the standard example  $u(x) = \log|\log(x)|$  shows that  $W^{1,p(\cdot)}(D) \not\hookrightarrow L^\infty$ , the calculations being as in the theorem. We next show that the exponent  $p$  from the theorem is almost as good as possible. We need the following lemma.

**Lemma 4.5.** *Let  $\{a_i\}$  be a partition of unity and  $k > m - 1$ . Then*

$$\sum_{i=0}^\infty a_i^m i^k \geq \left( \sum_{i=0}^\infty i^{-k/(m-1)} \right)^{1-m}.$$

*Proof.* Fix an integer  $i$  and consider the function

$$a \mapsto (a_i + a)^m i^k + (a_{i+1} - a)^m (i + 1)^k,$$

for  $-a_i < a < a_{i+1}$ . We find that this function has a minimum at  $a = 0$  if and only if

$$\left(\frac{a_i}{a_{i+1}}\right)^{m-1} = \left(\frac{i+1}{i}\right)^k. \tag{13}$$

Let  $\{a_i\}$  be a minimal sequence, so that (13) holds for every  $i \geq 0$ . This partition is given by  $a_i = i^{-k/(m-1)}a_0$  for  $i > 0$  and  $a_0 = (\sum i^{-k/(m-1)})^{-1}$  and so we easily calculate the lower bound as given in the lemma.  $\square$

We next give a simple sufficient condition for the imbedding  $W^{1,p(\cdot)}(D) \hookrightarrow L^\infty(D)$  to hold in a regular domain:

**Theorem 4.6.** *Suppose that  $D$  satisfies a uniform interior cone condition. If  $p^+ < \infty$  and*

$$p(x) \geq n + (n - 1 + \varepsilon) \frac{\log_2 \log_2(c/\delta(x))}{\log_2(c/\delta(x))}$$

for some fixed  $0 < \varepsilon < n - 1$  and constant  $c > 0$  then  $W^{1,p(\cdot)}(D) \hookrightarrow L^\infty(D)$ . Here  $\delta(x)$  denotes the distance of  $x$  from the boundary of  $D$

*Proof.* Note first that the claim trivially holds in compact subsets of  $D$  which satisfy the cone condition, since  $p$  is bounded away from  $n$  in such sets. Therefore it suffices to prove the claim for  $\delta(x)$  less than some constant.

By the uniform interior cone condition there exist real values  $0 < \alpha < \pi/2$  and  $r > 0$  and a unit vector field  $v_x$  such that for every  $x \in D$  the cone

$$C_x = \{y \in B(x, r) : \langle x - y, v_x \rangle > |x - y| \cos \alpha\}$$

lies completely in  $D$ , where  $\langle \cdot, \cdot \rangle$  denotes the usual inner product.

Fix  $z \in D$ . Consider the cone

$$C = \{y \in B(z, r/2) : \langle z - y, v_z \rangle > |z - y| \cos(\alpha/3)\}$$

and, for  $i = 2, 3, \dots$ , the annuli

$$A_i = (B(z, 2^{-i+1}r) \setminus B(z, 2^{-i}r)) \cap C.$$

To simplify notation let us assume that  $z = 0$ ,  $r = 1$  and  $v_z = e_1$ ; the proof in the general case is essentially identical. Since  $A_i \subset C \subset D$  we have  $d(A_i, \partial D) \geq d(A_i, \partial C)$ . We can estimate the latter distance as shown in Figure 1. This gives  $d(A_i, \partial D) \geq 2^{-i} \sin(\alpha/3)$  so that

$$p(x) \geq n + (n - 1 + \varepsilon) \frac{\log_2(i + c)}{i + c}$$




Figure 1: The cone  $C$  and the distance to the boundary

for  $x \in A_i$  and some  $c$  depending on  $\alpha$ . Let us define  $q_i = n + (n - 1 + \varepsilon) \frac{\log_2(i+c)}{i+c}$  and a new variable exponent by

$$q(x) = \begin{cases} q_i & \text{if } x \in A_i \text{ for some } i \\ p(x) & \text{otherwise} \end{cases}$$

By Theorem 4.3 we know that it suffices to find a lower bound for  $\|\nabla u\|_{1,p(\cdot)}$  with  $u \in L(\overline{D} \setminus B(0, r), 0; D)$  since, by Theorem 3.1,  $W^{1,p(\cdot)}(D) \subset C(D)$ . Since  $\|u\|_{1,p(\cdot)} \geq c\|u\|_{1,q(\cdot)}$ , we see that it suffices to estimate  $\psi_{q(\cdot)}(\overline{D} \setminus B(0, R), 0; B(0, R) \cap D)$  for small  $R$  in order to prove the theorem. Moreover, by monotony, we need only consider  $\psi_{q(\cdot)}(\overline{D} \setminus B(0, R), 0; B(0, R) \cap C)$ . For every function  $u \in W^{1,q(\cdot)}(C)$  we have

$$\|u\|_{1,q(\cdot)} \geq \min\{1, \varrho_{1,q(\cdot)}(u)\},$$

by [17, Theorem 2.8]. Therefore we see that it suffices to show that  $\varrho_{1,q(\cdot)}(u) > c$  for every  $u \in L(\overline{D} \setminus B(0, R), 0; B(0, R) \cap C)$  in order to get  $\psi_{q(\cdot)}(\overline{D} \setminus B(0, R), 0; B(0, R) \cap C) \geq \min\{1, c\} > 0$ , which will complete the proof.

It is clear that  $|\nabla u| \geq |\partial u / \partial r|$ , the radial component of the gradient, so that

$$\int_{A_i} |\nabla u|^{q_i} dx \geq \int_{A_i} \left| \frac{\partial u}{\partial r} \right|^{q_i} dx.$$

It is then easy to see that the function minimizing the sum over the integrals should depend only on the distance from the origin, not on the direction. For such a function let us denote the value at any point of distance  $2^{-i}$  from the origin by  $v_i$ .

Consider then a function  $v$  which equals  $v_{i-1}$  on  $S(0, 2^{-i+1})$  and  $v_i$  on  $S(0, 2^{-i})$ . Using Lemma 2.4 we find that

$$\begin{aligned} \int_{A_i} |\nabla v|^{q_i} dx &\geq (v_{i-1} - v_i)^{q_i} \text{cap}_{q_i}(\mathbb{R}^n \setminus B(0, 2^{-i+1}), B(0, 2^{-i})) \\ &= (v_{i-1} - v_i)^{q_i} \omega_{n-1} \left(\frac{q_i - n}{q_i - 1}\right)^{q_i-1} (2^{(q_i-n)/(q_i-1)} - 1)^{1-q_i} 2^{i(q_i-n)} \\ &\geq c(v_{i-1} - v_i)^{q_i} 2^{i(q_i-n)}, \end{aligned}$$

where the constant  $c$  does not depend on  $q_i$ . It follows that

$$\varrho_{1,q(\cdot)}(v) \geq \sum_{i=2}^{\infty} \int_{A_i} |\nabla v|^{q_i} dx \geq c \sum_{i=2}^{\infty} (v_{i-1} - v_i)^{q_i} 2^{i(q_i-n)}.$$

Since the lower bound depends only on the  $v_i$ , we see that

$$\inf_{u \in L} \varrho_{1,q(\cdot)}(u) \geq c \inf_{\{v_i\}} \sum_{i=2}^{\infty} (v_{i-1} - v_i)^{q_i} 2^{i(q_i-n)},$$

where the second infimum is over sequences  $\{v_i\}$  with  $v_i \leq v_{i-1}$ ,  $v_0 = 1$  and  $\lim_{i \rightarrow \infty} v_i = 0$ . Let us set  $a_i = v_{i-1} - v_i$  so that  $a_i \geq 0$  and  $\sum a_i = 1$ . Then we need to estimate

$$\inf_{\{a_i\}} \sum_{i=2}^{\infty} a_i^{q_i} 2^{i(q_i-n)},$$

with the infimum over partitions of unity  $\{a_i\}$ . Let  $N$  be such that

$$\frac{\varepsilon}{3} \geq q_i - n = (n - 1 + \varepsilon) \frac{\log_2(i + c)}{i + c} \geq (n - 1 + \varepsilon/2) \frac{\log_2(i)}{i}$$

for  $i \geq N$ . Note that such an  $N$  can be chosen independent of  $z$ . Since  $a_i \leq 1$  we have  $a_i^{q_i} \geq a_i^{n+\varepsilon/3}$  for such terms. Then we find that

$$\inf_{\{a_i\}} \sum_{i=2}^{\infty} a_i^{q_i} 2^{i(q_i-n)} \geq \inf_{\{a_i\}} \sum_{i=2}^{N-1} a_i^{q_i} 2^{i(q_i-n)} + \sum_{i=N}^{\infty} a_i^{n+\varepsilon/3} i^{n-1+\varepsilon/2}.$$

The first sum on the left-hand-side is finite, hence

$$\sum_{i=2}^{N-1} a_i^{q_i} 2^{i(q_i-n)} \geq \sum_{i=2}^{N-1} a_i^q \geq N^{1-q} \left(\sum_{i=2}^{N-1} a_i\right)^q,$$

where  $q = \max_{2 \leq i \leq N-1} q_i$ . It follows from Lemma 4.5 that

$$\sum_{i=N}^{\infty} a_i^{n+\varepsilon/3} i^{n-1+\varepsilon/2} \geq c \left(\sum_{i=N}^{\infty} a_i\right)^{n+\varepsilon/3}.$$



Combining these estimates we see that

$$\inf_{\{a_i\}} \sum_{i=2}^{\infty} a_i^{q_i} 2^{i(q_i-n)} \geq N^{1-q} \left( \sum_{i=2}^{N-1} a_i \right)^q + c \left( \sum_{i=N}^{\infty} a_i \right)^{n+\varepsilon/3}$$

is uniformly bounded from below by a positive constant, since the sum of the  $a_i$ 's is 1. We have thus shown that the condition of Theorem 4.3 holds, which concludes the proof.  $\square$

## References

- [1] S. M. Buckley and P. Koskela, *Sobolev-Poincaré implies John*, Math. Res. Lett. **2** (1995), 577–593.
- [2] ———, *Criteria for imbeddings of Sobolev-Poincaré type*, Internat. Math. Res. Notices (1996), 881–901.
- [3] B. Bojarski, *Remarks on Sobolev imbedding inequalities*, Complex Analysis, Joensuu 1987, Lecture Notes in Math., vol. 1351, Springer, Berlin, 1988, pp. 52–68.
- [4] D. Cruz-Uribe, A. Fiorenza, and C. J. Neugebauer, *The maximal function on variable  $L^p$  spaces*, Ann. Acad. Sci. Fenn. Math. **28** (2003), 223–238.
- [5] L. Diening, *Maximal function on generalized Lebesgue spaces  $L^{p(\cdot)}$* , Math. Inequal. Appl. (to appear).
- [6] ———, *Riesz Potential and Sobolev Embeddings of generalized Lebesgue and Sobolev Spaces  $L^{p(\cdot)}$  and  $W^{k,p(\cdot)}$* , Math. Nachr. (to appear).
- [7] L. Diening and M. Růžička, *Calderón-Zygmund operators on generalized Lebesgue spaces  $L^{p(\cdot)}$  and problems related to fluid dynamics*, J. Reine Angew. Math. **563** (2003), 197–220.
- [8] D. E. Edmunds and J. Rákosník, *Density of smooth functions in  $W^{k,p(x)}(\Omega)$* , Proc. Roy. Soc. London Ser. A **437** (1992), 229–236.
- [9] ———, *Sobolev embeddings with variable exponent*, Studia Math. **143** (2000), 267–293.
- [10] ———, *Sobolev embeddings with variable exponent. II*, Math. Nachr. **246/247** (2002), 53–67.
- [11] X. Fan, J. Shen, and D. Zhao, *Sobolev embedding theorems for spaces  $W^{k,p(x)}(\Omega)$* , J. Math. Anal. Appl. **262** (2001), 749–760.
- [12] P. Hajlasz and O. Martio, *Traces of Sobolev functions on fractal type sets and characterization of extension domains*, J. Funct. Anal. **143** (1997), 221–246.
- [13] P. Harjulehto, P. Hästö, M. Koskenoja, and S. Varonen, *Sobolev capacity on the space  $W^{1,p(\cdot)}(\mathbb{R}^n)$* , J. Funct. Spaces Appl. **1** (2003), 17–33.
- [14] ———, *Dirichlet energy integral and Sobolev spaces with zero boundary values* (preprint. Available in <http://www.math.helsinki.fi/analysis/varsobgroup>).
- [15] J. Heinonen, T. Kilpeläinen, and O. Martio, *Nonlinear potential theory of degenerate elliptic equations*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1993, ISBN 0-19-853669-0.
- [16] V. Kokilashvili and S. Samko, *Maximal and fractional operators in weighted  $L^{p(x)}$  spaces*, Rev. Mat. Iberoamericana (to appear).
- [17] O. Kováčik and J. Rákosník, *On spaces  $L^{p(x)}$  and  $W^{k,p(x)}$* , Czechoslovak Math. J. **41(116)** (1991), 592–618.

- [18] J. Malý and W. P. Ziemer, *Fine regularity of solutions of elliptic partial differential equations*, Mathematical Surveys and Monographs, vol. 51, American Mathematical Society, Providence, RI, 1997, ISBN 0-8218-0335-2.
- [19] O. Martio and J. Sarvas, *Injectivity theorems in plane and space*, Ann. Acad. Sci. Fenn. Ser. A I Math. **4** (1979), 383–401.
- [20] V. G. Maz'ja, *Sobolev spaces*, Springer Series in Soviet Mathematics, Springer-Verlag, Berlin, 1985, ISBN 3-540-13589-8.
- [21] L. Pick and M. Růžička, *An example of a space  $L^{p(x)}$  on which the Hardy-Littlewood maximal operator is not bounded*, Expo. Math. **19** (2001), 369–371.
- [22] Michael Růžička, *Electrorheological fluids: modeling and mathematical theory*, Lecture Notes in Mathematics, vol. 1748, Springer-Verlag, Berlin, 2000, ISBN 3-540-41385-5.