

# Tail and free poset algebras

Mohamed BEKKALI and Driss ZHANI

Faculty of Sciences and Technology

P.O.Box 2202

Fez 30000, Morocco

bekka@menara.ma

drisszhani@hotmail.com

Recibido: 4 de Septiembre de 2002

Aceptado: 22 de Octubre de 2003

*This work is dedicated to the  
philosopher Ibn Tophail (1100–1185).*

## ABSTRACT

We characterize free poset algebras  $F(P)$  over partially ordered sets and show that they can be represented by upper semi-lattice algebras. Hence, the uniqueness, in decomposition into normal form, using symmetric difference, of non-zero elements of  $F(P)$  is established. Moreover, a characterization of upper semi-lattice algebras that are isomorphic to free poset algebras is given in terms of a selected set of generators of  $B(T)$ .

*Key words:* Free poset algebra, tail algebra, semigroup algebra

*2000 Mathematics Subject Classification:* Primary: 03G, 06A; Secondary: 06A12, 03G10, 03G05

## 1. Characterization of $F(P)$

The main result of this section is Theorem 1.2 that gives an algebraic characterization of free poset algebras. Our approach in studying these algebras is different from [1] and [8]. Indeed, we show that the class of free poset algebras is actually a subclass of upper semi-lattice algebras. Therefore, non-zero elements has a unique decomposition into normal form. Two elements  $a$  and  $b$  of a poset  $\langle P, \leq \rangle$  are *comparable* whenever  $a \leq b$  or  $b \leq a$ . We say  $a \parallel b$ , whenever  $a$  and  $b$  are not comparable in  $\langle P, \leq \rangle$ . A subset  $A$  of  $\langle P, \leq \rangle$  of non comparable elements is called an *anti-chain* of  $\langle P, \leq \rangle$ .  $\mathcal{I}d(P)$  shall denote the set of ideals of  $P$ . ( $J \subseteq P$  is an ideal if  $J$  is a non empty initial segment of  $P$  such that every  $p, q \in J$  there is  $r \in J$  such that  $p, q \leq r$ ). For a non empty set  $X$ , we denote by  $[X]^{<\omega}$  the set all finite subsets of  $X$ . Thus, we set  $\text{Ant}(P) = \{\sigma \in [P]^{<\omega} : \sigma \text{ is an antichain of } P\}$ .

Now, a subset  $F$  of  $P$  is a *final segment* of  $P$ , whenever it is closed upwards i.e., if  $(a, b) \in F \times P$  and  $a \leq b$ , then  $b \in F$ . E.g., for each  $p \in P$ ,  $P^{\geq p} := \{q \in P : p \leq q\}$  is a final segment of  $P$ . For a finite subset  $\sigma$  of  $P$ ,  $P^{\geq \sigma} := \bigcup_{p \in \sigma} P^{\geq p}$  is a final segment of  $P$  generated by  $\sigma$ . Define  $\preceq$  be the binary relation on  $[P]^{<\omega}$  defined by  $\sigma \preceq \tau$  if  $P^{\geq \sigma} \supseteq P^{\geq \tau}$ . i. e., for every  $q \in \tau$  there is  $p \in \sigma$  such that  $p \leq q$ . So  $\preceq$  is a reflexive and transitive relation. For  $\sigma \in [P]^{<\omega}$  let  $\min(\sigma)$  be the set of minimal elements of  $\sigma$ . So  $P^{\geq \sigma} = P^{\geq \min(\sigma)}$ . It is trivial that  $\preceq$  restricted to  $\text{Ant}(P)$  is a partial order. Also for  $\sigma, \tau \in \text{Ant}(P)$ ,  $P^{\geq \sigma} \cup P^{\geq \tau} = P^{\geq \sigma \cup \tau} = P^{\geq \min(\sigma \cup \tau)}$ .

Throughout this paper  $\langle P, \leq \rangle$  shall denote a partially ordered set (*poset*) with greatest element  $\infty$ . Notice that this assumption on  $\langle P, \leq \rangle$  is not restrictive but it's rather convenient for expository purposes; indeed, let  $\langle Q, \leq \rangle$  be a poset and denote by  $\text{Fs}^+(Q)$  the set of all final segments of  $Q$ , and  $\text{Fs}(Q) := \text{Fs}^+(Q) \setminus \{\emptyset\}$  then  $\text{Fs}(Q)$  and  $\text{Fs}(Q \cup \{\infty\})$  are homeomorphic spaces; for more see e. g. [3].  $\text{Fs}(P)$  the set of all non-empty final segments of  $P$  is a closed subspace of  $\{0, 1\}^P$  by identifying  $\wp(P)$  with  $\{0, 1\}^P$  via characteristic functions of sets and taking the the Cantor topology on  $\{0, 1\}^P$ . For a topological space  $X$ ,  $\text{clop}(X)$  shall denote the Boolean algebra of clopen (*closed and open*) subsets of  $X$ . Thus,  $\text{clop}(\text{Fs}(P))$  is the *free poset algebra* over  $P$ , denoted by  $F(P)$ . For  $p \in P$ , set  $V_p := \{F \in \text{Fs}(P) : p \in F\}$ . So  $V_p$  is a clopen subset of  $\text{Fs}(P)$ . Also, for any finite subset  $\sigma$  of  $P$ .  $(\dagger) \bigcap_{p \in \sigma} V_p = \{F \in \text{Fs}(P) : F \supseteq P^{\geq \sigma}\}$ . Hence a basis of  $\text{Fs}(P)$  is  $\langle \bigcap_{p \in \sigma} V_p \cap \bigcap_{q \in \tau} -V_q, \sigma, \tau \in [P]^{<\omega} \rangle$ , where  $-V_q$  denotes the complement of  $V_q$  in  $\text{Fs}(P)$ ; (i.e.,  $-V_q = \text{Fs}(P) \setminus V_q$ ). Notice that  $V_p \in F(P)$  and that every member of  $F(P)$  is a finite union of  $\bigcap_{p \in \sigma} V_p \cap \bigcap_{q \in \tau} -V_q$ . Finally, recall that  $\text{cl}_B(S)$  denotes the subalgebra of  $B$  generated by  $S$  for any Boolean algebra  $B$  and any  $S \subseteq B$ . Next,  $\text{Ult}(B)$ , the set of ultrafilters of  $B$ , shall denote the Stone space of  $B$ .

**Lemma 1.1.** *Let  $\langle P, \leq \rangle$  be a poset, with greatest element  $\infty$ .*

- (i) *If  $p < q$  then  $V_p \subseteq V_q$ .*
- (ii)  $\bigcap_{p \in P} V_p = \{P\} \neq \emptyset$ .
- (iii)  $V_\infty = \text{Fs}(P)$ .
- (iv) *For finite subsets  $\sigma, \tau$  of  $P$ , the following properties are equivalent:*
  - (a)  $\bigcap_{p \in \sigma} V_p \cap \bigcap_{q \in \tau} -V_q = \emptyset$ .
  - (b) *There are  $p \in \sigma, q \in \tau$  such that  $p \leq q$ .*
- (v) *If  $\bigcap_{i=1}^n V_{p(i)} \subseteq \bigcup_{k=1}^m \left( \bigcap_{j=1}^{m(k)} V_{q(k,j)} \right)$  then*
  - (a) *there is  $k \in \{1, \dots, m\}$ , such that  $\bigcap_{i=1}^n V_{p(i)} \subseteq \bigcap_{j=1}^{m(k)} V_{q(k,j)}$ , and*
  - (b) *for any  $j \in \{1, \dots, m(k)\}$  there is  $i \in \{1, \dots, n\}$  so that  $p(i) \leq q(k, j)$ .*

*Proof.* (i)–(iii) are easy to check.

To prove (iv), set  $W := \bigcap_{p \in \sigma} V_p \cap \bigcap_{q \in \tau} -V_q$ . Then,

$$W = \{F \in \text{Fs}(P) : \sigma \subseteq F \text{ and } \tau \cap F = \emptyset\}.$$

Suppose that there are  $p \in \sigma, q \in \tau$  and  $p \leq q$ . So every final segment containing  $\sigma$  contains  $q$ . Thus,  $W = \emptyset$ . Conversely, suppose that for every  $p \in \sigma, q \in \tau$ , and  $p \not\leq q$  we have  $P^{\geq \sigma} \cap \tau = \emptyset$ . It follows then,  $P^{\geq \sigma} \in W$ ; thus  $W \neq \emptyset$ .

To prove (v), set  $\sigma := \{p(1), \dots, p(n)\}, F := P^{\geq \sigma}$ . So,  $F \in \bigcap_{i=1}^n V_{p(i)}$ . Choose  $k$  so that  $F \in \bigcap_{j=1}^{m(k)} V_{q(k,j)}$ . Note that  $q(k, j) \in F$  for every  $j \in \{1, \dots, m(k)\}$ . Now, if  $G \in \bigcap_{i=1}^n V_{p(i)}$  then  $F \subseteq G$ ; thus  $q(k, j) \in G$  for every  $j \in \{1, \dots, m(k)\}$ . Hence,  $\bigcap_{i=1}^n V_{p(i)} \subseteq \bigcap_{j=1}^{m(k)} V_{q(k,j)}$ . Next, since  $F \in \bigcap_{j=1}^{m(k)} V_{q(k,j)}$  it follows that for every  $q(k, j)$  there is  $p(i)$  such that  $p(i) \leq q(k, j)$ .  $\square$

The following theorem characterizes free poset algebras and will be of use later on in the paper.

**Theorem 1.2.** *The following statements are equivalent for any Boolean algebra  $B$ .*

- (i)  $B$  is isomorphic to a free poset algebra.
- (ii)  $B$  has a set  $H$  of generators with  $1 \in H$  such that for every finite subsets  $\{h_i : i < m\}$  and  $\{k_j : j < n\}$  of  $H$ :

(a)  $\prod_{i < m} h_i \neq 0$  and

(b) if  $\prod_{i < m} h_i \cdot \prod_{j < n} -k_j = 0$  then there are  $i$  and  $j$  such that  $h_i \leq k_j$ .

*Proof.* (i) implies (ii). We may assume that  $B = F(P)$ . Let  $H = \{V_q : q \in P\}$ . By Lemma 1.1, (ii) holds.

(ii) implies (i). Let  $H$  be as in (ii). So  $H$  is a poset with a greatest element. To show that  $B$  is isomorphic to  $F(H)$ , let  $f : H \rightarrow F(H)$  be defined by  $f(h) = V_h$ . By Lemma 1.1 (iv) and the hypothesis (ii)-(b), the following are equivalent:

1.  $\prod_{i < m} h_i \cdot \prod_{j < n} -k_j = 0$
2. there is  $i$  and  $j$  such that  $h_i \leq k_j$ , and
3.  $\bigcap_{i < m} V_{h_i} \cap \bigcap_{j < n} -V_{k_j} = \emptyset$

By Sikorski's Criterion,  $f$  extends to an isomorphism  $\hat{f}$  from  $\text{cl}_B(H)$  onto  $\text{Im}(f) \subseteq F(H)$ . Since  $H$  generates  $B$ , and  $\text{Im}(f) = F(H)$ ,  $\hat{f}$  is an isomorphism from  $B$  onto  $F(H)$ .  $\square$

## 2. Representation of $F(P)$

Let  $\langle Q, \leq \rangle$ . For  $q \in Q$ , put  $b_q := \{u \in Q : u \geq q\}$ . Next define the *Tail algebra*,  $B(Q)$ , as the subalgebra of the power set of  $Q$  generated by  $\{b_q : q \in Q\}$ . When  $\langle T, \leq \rangle$  is an *upper semi-lattice* i. e., l. u. b.  $\{x, y\} := x \vee y$  exists in  $\langle T, \leq \rangle$  for  $x, y \in T$ ;  $B(T)$  is called the *upper semi-lattice algebra* over  $T$ . Notice that every member of  $B(T)$  is a finite union of  $\bigcap_{t \in \sigma} b_t \cap \bigcap_{s \in \tau} -b_s$  (where  $\sigma, \tau$  are finite subsets of  $T$ ).

In this section Theorem 2.3 shows that any  $F(P)$  is isomorphic to an upper semi-lattice algebra. As for Theorem 2.4, a characterization of upper semi-lattice algebras that are isomorphic to an  $F(P)$  is given using the idea of *prime elements* in the upper semi-lattice.

Before we state the following lemmas, notice that if  $Q$  is the poset  $\{p, q, r, s\}$  with relations  $p < r < q$ ,  $p < s < q$  and  $r, s$  incomparable; then in the tail algebra  $B(Q)$ ,  $b_r \cap b_s = b_q$ ; nevertheless, in the free poset algebra  $F(Q)$ ,  $V_p \subset V_r \cap V_s$  and  $V_r \cup V_s \subset V_q$ .

**Lemma 2.1.** *Let  $B(Q)$  be the tail algebra over  $Q$ . Let  $p \in Q$  and  $\tau$  be a finite subset of  $Q$ . The following properties are equivalent.*

- (i)  $b_p \subseteq \bigcup_{q \in \tau} b_q$ .
- (ii)  $q \in \tau$  such that  $q \leq p$ .

**Lemma 2.2.** (i)  $\preceq$  is an ordering on  $\text{Ant}(P)$ .

- (ii) The greatest lower bound of  $\sigma$  and  $\tau$  is  $\min(\sigma \cup \tau)$  in  $\langle \text{Ant}(P), \preceq \rangle$ , for  $\sigma, \tau \in \text{Ant}(P)$ ,

- (iii)  $\langle \text{Ant}(P), \succeq \rangle$  is an upper semi-lattice, with a least element.

Next theorem shows that the class of upper semi-lattice algebras contains the class of free poset algebras.

**Theorem 2.3.** *For every poset  $(P, \leq)$ , with a greatest element,  $F(P)$  is isomorphic to the upper semi-lattice algebra  $B(\langle \text{Ant}(P), \succeq \rangle)$ .*

*Proof.* Set  $H = \{V_q : q \in P\}$ . Recall that  $H$  generates  $F(P)$ . Now, for each  $p \in P$ ,  $\{p\} \in \text{Ant}(P)$ ; and thus  $b_{\{p\}} = \{\sigma \in \text{Ant}(P) : \sigma \preceq \{p\}\}$ .

Next, define  $\varphi : H \rightarrow B(\langle \text{Ant}(P), \succeq \rangle)$  by  $\varphi(V_p) = b_{\{p\}}$ . We claim that for every  $\sigma, \tau \in \text{Ant}(P)$ :

$$\bigcap_{p \in \sigma} V_p \cap \bigcap_{q \in \tau} -V_q = \emptyset \quad \text{iff} \quad \bigcap_{p \in \sigma} b_{\{p\}} \cap \bigcap_{q \in \tau} -b_{\{q\}} = \emptyset. \tag{1}$$

By Lemma 1.1 (iv), we have:

$$\bigcap_{p \in \sigma} V_p \cap \bigcap_{q \in \tau} -V_q = \emptyset \quad \text{iff} \quad \text{“there are } p \in \sigma \text{ and } q \in \tau \text{ so that } p \leq q\text{”} \quad (2)$$

On the other hand, we have  $\{\{p\} : p \in \sigma\} \subseteq \text{Ant}(P)$  and  $\min(\sigma)$  is its l.u.b. in  $\langle \text{Ant}(P), \succeq \rangle$ . So  $\bigcap_{p \in \sigma} b_{\{p\}} = b_{\min(\sigma)}$ . So  $\bigcap_{p \in \sigma} b_{\{p\}} \cap \bigcap_{q \in \tau} -b_{\{q\}} = \emptyset$ ; which means that  $b_{\min(\sigma)} \subseteq \bigcup_{q \in \tau} b_{\{q\}}$ . By Lemma 2.1, this is equivalent to the existence of  $q \in \tau$  so that  $\{q\} \succeq \min(\sigma)$ .

Next  $\{q\} \succeq \sigma$  iff there is  $p \in \sigma$  so that  $p \leq q$ , that is (2). The proof of (1) is finished.

Next, by (1) and Sikorski’s Criterion,  $\varphi$  extends to a monomorphism  $\hat{\varphi}$  from  $F(\langle P, \leq \rangle)$  into  $B(\langle \text{Ant}(P), \succeq \rangle)$ . Now, since  $\{V_p : p \in P\}$  generates  $F(P)$ ,  $\{b_{\{p\}} : p \in P\}$  generates  $B(\text{Ant}(P))$  and  $\hat{\varphi}(V_p) = b_{\{p\}}$ ; the isomorphism is established and the proof of the theorem is finished.  $\square$

Recall that an element  $p \in T$  is called a *prime* element of  $T$  whenever for every  $u, v \in T$  so that  $u \vee v$  exists in  $T$  and  $p \leq u \vee v$ , then  $p \leq u$  or  $p \leq v$ .  $\text{Prim}(T)$  shall denote the set of all prime elements of  $T$ .

Next is a characterization of free poset algebras.

**Theorem 2.4.** *Let  $T$  be an upper semi-lattice. The following statement are equivalent.*

- (i)  $B(T)$  is isomorphic to a free poset algebra.
- (ii) There is a poset  $P$ , with a greatest element so that  $B(T) \cong B(\langle \text{Ant}(P), \succeq \rangle)$ .
- (iii) There is an upper semi-lattice  $T'$ , with a least element, so that:
  - (a) Every element of  $T'$  is a join of finitely many prime elements,
  - (b)  $B(T)$  and  $B(T')$  are isomorphic Boolean algebras, and
  - (c)  $B(T') = \text{cl}_{B(T')}(\{b_t : t \in \text{Prim}(T')\})$ .

*Proof.* (i) implies (ii) follows from Theorem 2.3.

(ii) implies (iii). Let  $\langle T', \leq \rangle := \langle \text{Ant}(P), \succeq \rangle$ . By Lemma 2.2 (ii),  $T'$  is an upper semi-lattice. Next, it is straightforward to notice that  $\sigma \in \text{Ant}(P)$ :  $\sigma$  is prime in  $\langle \text{Ant}(P), \succeq \rangle$  whenever  $\sigma$  is a singleton. Note that by the proof of Theorem 2.3,  $\{b_{\{p\}} : p \in P\}$  generates  $B(\text{Ant}(P), \succeq)$ .

(iii) implies (i). Let  $T'$  be as in (iii). Let  $t_0$  be the least element of  $T'$ . Let  $P = \text{Prim}(T')$ . (So  $P \subseteq T'$ .) Let  $H' := \{b_t : t \in P\}$ . By (iii)-(c)

$$B(T') = \text{cl}_{B(T')}(H'). \quad (3)$$

Since  $t_0 \in P$ :

$$1^{B(T')} = b_{t_0} \in H'. \quad (4)$$

For  $\{t(1), \dots, t(n)\} \subseteq P$ ,  $\bigcap_{i=1}^n b_{t(i)} = b_{t(1) \vee \dots \vee t(n)} \neq 0$ , and thus:

$$H \text{ has the finite intersection property.} \tag{5}$$

For  $\{t(1), \dots, t(m)\} \subseteq P$  and  $\{s(1), \dots, s(n)\} \subseteq P$ . we have:

$$\bigcap_{i=1}^m b_{t(i)} \cap \bigcap_{j=1}^n -b_{s(j)} = \emptyset \quad \text{whenever there are } i \text{ and } j \text{ so that } s(j) \leq t(i). \tag{6}$$

To see that (6) holds set  $t = \bigvee_i t(i)$ . So the left hand side of (6) is equivalent to  $b_t \subseteq \bigcup_{j=1}^n b_{s(j)}$ . In other words,  $t \in \bigcup_{j=1}^n b_{s(j)}$ . i. e.,  $t \geq s(j)$  for some  $j$ . Since  $s(j) \in P = \text{Prim}(T')$ , this is equivalent to say that there is  $i$  so that  $t(i) \geq s(j)$ .

Next, let  $H := \{V_p : p \in P\} \subseteq F(P)$  and  $f : H' \rightarrow H$  defined by  $f(b_p) = V_p$ . By (6), Lemma 1.1 (iv) and using Sikorski's Criterion,  $f$  extends to a monomorphism  $\hat{f}$  from  $\text{cl}_{B(T')}(H')$  into  $F(P)$ . By (3),  $\text{Dom}(f) = B(T')$ , and since  $H$  generates  $F(P)$ ,  $\hat{f}$  is actually onto and thus, an isomorphism.  $\square$

The following proposition summarizes the relationship between  $T$  and  $P$  whenever  $B(T) \cong F(P)$ .

- Proposition 2.5.** (i) *If a poset  $\langle P, \leq \rangle$  with a greatest element is so that  $F(P) \cong B(T)$ , then  $T$  may be chosen canonically to be  $\langle T, \leq \rangle = \langle \text{Ant}(P), \succeq \rangle$ .*
- (ii) *If an upper semi-lattice  $T$  is so that  $B(T) \cong F(P)$ , then an upper semi-lattice  $T'$  may be chosen so that  $B(T) \cong B(T') = \text{cl}_{B(T')}(\{b_t : t \in \text{Prim}(T')\})$  and  $P \cong \{b_t : t \in \text{Prim}(T')\}$ .*

### 3. Normal form of non-zero elements in the free poset algebra $F(P)$

In this section we show, by Lemma 3.4, that any non-zero element of  $F(P)$  has a decomposition into normal form as it is the case in the class of pseudo-tree algebras see, for instance, [2] and [6, p. 51]. Unfortunately, this representation of elements, here in the free poset algebra  $F(P)$ , is not unique. Using symmetric difference  $\Delta$ , instead, we shall see that Theorem 3.5 gives uniqueness in normal form of non-zero elements.

Before we give next lemma, recall that if  $P$  is a poset then  $\langle \text{Fs}(P), \subseteq \rangle$  is a poset too. Thus, we consider the tail algebra over  $\langle \text{Fs}(P), \subseteq \rangle$ . For notational purposes, we use small letters for elements of  $\text{Fs}(P)$ , and  $\leq$  denotes the inclusion relation. Thus, for  $p \in P$ , we set  $h_p = P^{\geq p}$ . Hence, in the tail algebra  $B(\langle \text{Fs}(P), \subseteq \rangle)$ :

$$b_{h_p} = \{x \in \text{Fs}(P) : x \geq h_p\}$$

The following Lemmas follow easily.

**Lemma 3.1.** *Let  $P$  be a poset and  $q \in P$ . We have  $V_q = b_{h_q}$ .*

**Lemma 3.2.** *Let  $P$  be a poset and  $p, q \in P$ .*

- (i)  $p \leq q$  iff  $h_q \leq h_p$ .
- (ii)  $p \parallel q$  iff  $h_q \parallel h_p$ .
- (iii) For each  $q \in P$ ,  $h_q \in \text{Prim}(\text{Fs}(P))$ .
- (iv)  $\bigcap_{i \in \sigma} V_{q(i)} = \bigcap_{i \in \sigma} b_{h_{q(i)}} = b_{\bigvee_{i \in \sigma} h_{q(i)}}$  for each finite set  $\sigma$ .

Let  $P$  be a poset. Let

$$T^P := \left\{ \bigvee_{i \in \sigma} h_{q(i)} : \{q(i) : i \in \sigma\} \in \text{Ant}(P) \right\}.$$

Before we state next Lemma, we denote  $T^P$  by  $T$ .

**Lemma 3.3.** *Assume that  $\langle P, \leq \rangle$  is a poset with a greatest element  $\infty$ . Then:*

- (i)  $V_\infty = \text{Fs}(P)$ ;
- (ii)  $\langle T, \leq \rangle$  is an upper semi-lattice with a least element  $h_\infty$ ;
- (iii) For each  $f \in T$ ,  $-b_f = b_{h_\infty} \cdot -b_f$  and  $h_\infty \leq f$ ;
- (iv) For all  $f, g \in T$  we have:
  - (a)  $f \leq g$  iff  $b_g \cdot -b_f = \emptyset$ .
  - (b) If  $f \not\leq g$ , then  $b_g \cdot -b_f = b_g \cdot -b_{f \vee g}$ .

Next, we prove the first lemma concerning the decomposition of non zero elements in the free Boolean algebra  $F(P)$ .

**Lemma 3.4 (First normal form).** *Every  $b \in F(P) \setminus \{0\}$  can be written as  $b = e_1 + \dots + e_n$  where  $e_i \cdot e_j = 0$  for  $i \neq j$ , and for every  $i \in \{1, \dots, n\}$ , either  $e_i = b_{h_i}$ , or there is a finite anti-chain  $\{f_1, \dots, f_m\}$  in  $\langle T, \leq \rangle$  such that  $e_i = b_{h_i} \cdot -(b_{f_1} + \dots + b_{f_m})$ .*

*Proof.* Working out the proof, as in Proposition 4.4 in [6, p. 51], it suffices to show that each elementary product  $\prod_{i=1}^n \varepsilon_i V_{q_i}$  can be written, in  $F(P)$ , under the form  $b_h \cdot -(b_{f_1} + \dots + b_{f_m})$  with  $h, f_i \in T$ ,  $h < f_i$ , and  $\{f_1, \dots, f_m\}$  is an anti-chain.

Note that

$$\prod_{i=1}^n V_{q_i} = \bigcap_{i=1}^n V_{q_i} = \bigcap_{i=1}^n b_{h_{q_i}} = b_{\bigvee_{1 \leq i \leq n} h_{q_i}}$$

and that, by Lemma 3.3 (iii),

$$\prod_{i=1}^n -V_{q_i} = \bigcap_{i=1}^n (-b_{h_{q_i}}) = \bigcap_{i=1}^n (b_{h_\infty} \cdot -b_{h_{q_i}}) = b_{h_\infty} \cdot -(b_{h_{q_1}} + \dots + b_{h_{q_n}})$$

So we may assume that there are  $i, j$  so that  $\varepsilon_i = 1$  and  $\varepsilon_j = -1$ .

Thus  $\prod_{i=1}^n \varepsilon_i V_{q_i}$  can be written as,

$$\begin{aligned} \prod_{i=1}^n \varepsilon_i V_{q_i} &= V_{\alpha(1)} \cdots V_{\alpha(k)} \cdots -V_{\beta(1)} \cdots -V_{\beta(l)} \\ &= b_{h_{\alpha(1)}} \cdots b_{h_{\alpha(k)}} \cdots -b_{h_{\beta(1)}} \cdots -b_{h_{\beta(l)}} \\ &= b_{\vee_{1 \leq i \leq k} h_{\alpha(i)}} \cdots -b_{h_{\beta(1)}} \cdots -b_{h_{\beta(l)}} \end{aligned}$$

Now, set  $h = \vee_{1 \leq i \leq k} h_i$ . So,

$$\begin{aligned} \prod_{i=1}^n \varepsilon_i V_{q_i} &= b_h \cdots -b_{h_{\beta(1)}} \cdots -b_{h_{\beta(l)}} \\ &= (b_h \cdots -b_{h_{\beta(1)}}) \cdots (b_h \cdots -b_{h_{\beta(2)}}) \cdots (b_h \cdots -b_{h_{\beta(l)}}) \end{aligned}$$

Since  $\prod_{i=1}^n \varepsilon_i V_{q_i} \neq 0$ , by Lemma 3.3 (iv)-(b), for each  $j \in \{1, \dots, l\}$  either  $h < h_{\beta(j)}$  or  $h \parallel h_{\beta(j)}$ ; moreover  $b_h \cdots -b_{h_{\beta(j)}} = b_h \cdots -b_{f(j)}$  with  $h < f(j)$  and  $f(j) \in T$ . So,

$$\prod_{i=1}^n \varepsilon_i V_{q_i} = (b_h \cdots -b_{f(1)}) \cdots (b_h \cdots -b_{f(l)}) = b_h \cdots -(b_{f(1)} + \cdots + b_{f(l)})$$

Now, by canceling some of  $b_{f(i)}$ 's, if necessary, we may write

$$\prod_{i=1}^n \varepsilon_i V_{q_i} = b_h \cdots -(b_{f(i(1))} + \cdots + b_{f(i(m))})$$

where  $\{f(i(1)), \dots, f(i(m))\}$  is an anti-chain and  $h < f(i(k))$  for each  $k$ . This finishes up the proof of Lemma 3.4. □

*Remark.* Notice that normal form of non zero elements of  $F(P)$ , given by Lemma 3.4, may not be unique as shown by the following counterexample.

Let  $p, q \in P$  with  $p \parallel q$  and set  $b = b_{h_p} + b_{h_q}$ . we have

$$b = \underbrace{b_{h_p} \cdots -b_{h_p \vee h_q}}_{e_1} + \underbrace{b_{h_q}}_{e_2} = \underbrace{b_{h_p}}_{e'_1} + \underbrace{b_{h_q} \cdots -b_{h_p \vee h_q}}_{e'_2}$$

Indeed, by Lemma 3.3 (iv)-(b),  $b_{h_p} \cdots -b_{h_p \vee h_q} = b_{h_p} \cdots -b_{h_q}$ . Thus,

$$b_{h_p} \cdots -b_{h_p \vee h_q} + b_{h_q} = b_{h_p} \cdots -b_{h_q} + b_{h_q} = b_{h_p} + b_{h_q} = b.$$

Next, before we state the main theorem in this section, recall that whenever  $P$  is a poset,  $T := \{\vee_{i \in \sigma} h_{q(i)} : \{q(i) : i \in \sigma\} \in \text{Ant}(P)\}$  will be the upper semi-lattice that is going to be referred to in the next theorem.

**Theorem 3.5 (Normal form of non-zero elements in  $F(\mathbf{P})$ ).** *Every  $b \in F(\mathbf{P}) \setminus \{0\}$  has a unique decomposition as  $b = b_{g_1} \Delta \cdots \Delta b_{g_n}$ , where  $g_i \in T$ , and  $g_i \neq g_j$  for  $i \neq j$ .*

The proof of this theorem uses the following lemma.

**Lemma 3.6.** (i) *For all  $f, h \in T$ ,  $b_h \cdot -b_f = b_h \Delta b_{f \vee h}$ ,*

(ii) *If  $b = b_{f_1} \Delta \cdots \Delta b_{f_n}$  then  $b \cdot -b_f = \Delta_{i=1}^m b_{g_i}$  where  $g_i \in T$ .*

(iii) *If  $h < f_i$  (for every  $i$ ), then  $b_h \cdot -(b_{f_1} + \cdots + b_{f_n}) = \Delta_{i=1}^m b_{g_i}$  —where  $g_i \in T$ .*

(iv) *Set  $\min \{f_1, \dots, f_n\} = \{f_{i(1)}, \dots, f_{i(p)}\}$ . Then,*

$$b_{f_1} \Delta \cdots \Delta b_{f_n} \subseteq b_{f_{i(1)}} \cup \cdots \cup b_{f_{i(p)}} \quad \text{and} \quad f_{i(k)} \in b_{f_1} \Delta \cdots \Delta b_{f_n}.$$

(v) *If  $b_{f_1} \Delta \cdots \Delta b_{f_n} = b_{g_1} \Delta \cdots \Delta b_{g_m} \neq 0$ , then there are  $i, j$  so that  $b_{f_i} = b_{g_j}$ .*

*Proof.* (i) If  $f \leq h$  then  $b_h \cdot -b_f = \emptyset = b_h \Delta b_h$ . If  $f \not\leq h$ , by Lemma 3.3 (iv)-(b) we have,  $b_h \cdot -b_f = b_h \cdot -b_{f \vee h}$  and,

$$b_h \Delta b_{f \vee h} = b_h \cdot -b_{f \vee h} + \underbrace{b_{f \vee h} \cdot -b_h}_{=\emptyset} = b_h \cdot -b_{f \vee h}$$

Thus,  $b_h \cdot -b_f = b_h \Delta b_{f \vee h}$ .

(ii) Let

$$\begin{aligned} b &= b_{f_1} \Delta \cdots \Delta b_{f_n} \\ b \cdot -b_f &= (b_{f_1} \Delta \cdots \Delta b_{f_n}) \cdot -b_f \\ &= (b_{f_1} \cdot -b_f) \Delta \cdots \Delta (b_{f_n} \cdot -b_f) \\ &= (b_{f_1} \Delta b_{f \vee f_1}) \Delta \cdots \Delta (b_{f_n} \Delta b_{f \vee f_n}) \end{aligned}$$

(iii) Let  $h < f_i$  and show by induction on  $n$  that:

$$b_h \cdot -\sum_{i=1}^n b_{f_i} = \Delta_{i=1}^m b_{g_i}$$

Suppose that we have shown what we wanted up to  $n - 1$ . Let  $b = b_h \cdot -(b_{f_1} + \cdots + b_{f_{n-1}} + b_{f_n})$  and set  $b' = b_h \cdot -(b_{f_1} + \cdots + b_{f_{n-1}})$ . So,  $b = b_h \cdot -b_{f_1} \cdots \cdots -b_{f_{n-1}} \cdot -b_{f_n} = b' \cdot -b_{f_n}$ . Now, by induction hypothesis  $b' = \Delta_{i=1}^k b_{g_i}$ . So, by (ii),  $b = b' \cdot -b_{f_n} = (\Delta_{i=1}^k b_{g_i}) \cdot -b_{f_n} = \Delta_{j=1}^m b_{h_j}$ .

(iv) If  $f < g$  then  $b_g \subseteq b_f$ . So,

$$b_{f_1} \Delta \cdots \Delta b_{f_n} \subseteq b_{f_1} \cup \cdots \cup b_{f_n} \subseteq b_{f_{i(1)}} \cup \cdots \cup b_{f_{i(p)}}.$$

For each  $k \in \{1, \dots, p\}$ ,  $f_{i(k)} \in b_{f_1} \Delta \cdots \Delta b_{f_n}$ . Indeed,  $f_{i(k)} \in b_{f_{i(k)}}$  and for  $j \neq i(k)$ ,  $f_{i(k)} \notin b_{f_j}$  (if not  $f_j < f_{i(k)}$ ). So,

$$f_{i(k)} \in b_{f_{i(k)}} \Delta (\Delta_{j \neq i(k)} b_{f_j}) = b_{f_1} \Delta \cdots \Delta b_{f_n}.$$

(v) Suppose  $b := b_{f_1} \Delta \cdots \Delta b_{f_n} = b_{g_1} \Delta \cdots \Delta b_{g_m}$ , where  $f_i \neq f_j$  and  $g_k \neq g_l$  for  $i \neq j$  and  $k \neq l$ . Let

$$\min \{f_1, \dots, f_n\} = \{f_{i(1)}, \dots, f_{i(p)}\}, \quad \min \{g_1, \dots, g_m\} = \{g_{j(1)}, \dots, g_{j(q)}\}$$

By (iv) we have:

$$b \subseteq b_{f_{i(1)}} \cup \cdots \cup b_{f_{i(p)}} \tag{7}$$

$$b \subseteq b_{g_{j(1)}} \cup \cdots \cup b_{g_{j(q)}} \tag{8}$$

$$\text{for all } k \text{ and } \ell, \quad f_{i(k)} \in b \text{ and } g_{j(\ell)} \in b. \tag{9}$$

Let  $k$  be given. So, by (9)  $f_{i(k)} \in b$ . By (8), let  $\ell(k)$  be such that  $f_{i(k)} \geq g_{j(\ell(k))}$ . Similarly (using (7) instead of (8)), let  $k'$  be such that  $g_{j(\ell(k))} \geq f_{i(k')}$ . Hence  $f_{i(k)} \geq f_{i(k')}$ , and thus  $f_{i(k)} = f_{i(k')} = g_{j(\ell(k))}$ .  $\square$

Now we prove Theorem 3.5.

*Proof of Theorem 3.5.* We prove first the existence. Let  $b \in F(P)$ . By Lemma 3.4,  $b = \sum_{i=1}^n e_i$  where  $e_i \cdot e_j = \emptyset$  for  $i \neq j$  and thus  $b = \Delta_{i=1}^n e_i$ . In addition, by Lemma 3.4 again, either  $e_i = b_{h_i}$ , or  $e_i = b_{h_i} \cdot -(\sum_{j=1}^n b_{f_j})$ , and by Lemma 3.6 (iii)  $e_i = \Delta_{i=1}^m b_{g_i}$  ( $g_i \in T$ ).

We prove the uniqueness. Let  $b = \Delta_{i=1}^n b_{f_i} = \Delta_{j=1}^m b_{g_j} \neq 0$ . By Lemma 3.6(v) there are  $i, j$  so that  $b_{f_i} = b_{g_j}$ . Without loss of generality,  $i = 1 = j$ . So  $f_1 = g_1 := h$ . We have  $b_h \Delta b = \Delta_{i=2}^n b_{f_i} = \Delta_{j=2}^m b_{g_j}$ . The uniqueness follows.  $\square$

### 4. Examples of free poset algebras $F(P)$

The following proposition characterizes atoms in  $F(P)$ . To this end, let  $Atom(F(P))$  denotes the set of atoms of  $F(P)$ , and recall that, in  $F(P) = \langle V_q : q \in P \rangle$ , each element of  $F(P)$  is a finite union of  $\bigcap_{p \in \sigma} V_p \cap \bigcap_{q \in \tau} -V_q$ . The proof of the next result is obvious.

**Proposition 4.1.** *Let  $P$  be a poset. The following properties are equivalent.*

- (i)  $b \in Atom(F(P))$ .
- (ii) *There are finite subsets  $\sigma$  and  $\tau$  of  $P$  such that  $b = \bigcap_{p \in \sigma} V_p \cap \bigcap_{q \in \tau} -V_q$  and for every  $s \in P$ , either there is  $p \in \sigma$  such that  $s \geq p$  or there is  $q \in \tau$  such that  $s \leq q$ .*

Let us give some examples.

1. Let  $C_1, C_2$  be two chains with least elements  $\alpha, \beta$  respectively. Then  $B(C_1 \times C_2) \cong F(P)$ , for some poset  $P$ . Indeed,

$$Prim(C_1 \times C_2) = \{(\alpha, q) : q \in C_2\} \cup \{(p, \beta) : p \in C_1\}.$$

Moreover  $b_{(s,t)} = b_{(\alpha,t)} \vee b_{(s,\beta)}$ . Thus  $B(C_1 \times C_2) = \text{cl}(\{b_{(s,t)} : (s,t) \in \text{Prim}(C_1 \times C_2)\})$ . Now by Theorem 2.4 (iii),  $B(C_1 \times C_2) \cong F(P)$ , for some poset  $P$ .

2. For every set  $I$ , there is a poset  $\langle P, \leq \rangle$  with a greatest element so that the free Boolean algebra over  $I$  is isomorphic to  $F(P)$ . (Consider  $I$  as an anti-chain and  $P = I \cup \{\infty\}$  with  $p < \infty$  for every  $p \in I$ .)

The following proposition shows that free poset algebras is a proper subclass of upper semi-lattice algebras.

**Proposition 4.2.** *Let  $T$  be an anti-chain of size  $\aleph_1$  so that  $x \vee y =_{\text{def}} \infty$  for all  $x, y \in T$ . Then  $B(T \cup \{\infty\})$  is an upper semi-lattice algebra that is not a free poset algebra.*

*Proof.* Suppose the contrary and pick  $P$  so that  $\mathcal{I}d(T \cup \{\infty\}) = \text{Ult}(B(T \cup \{\infty\}))$  and  $\text{Fs}(P)$  are homeomorphic spaces. It follows that  $\text{Fs}(P)$  is a scattered topological space and thus  $|\text{Fs}(P)| = |P| = \aleph_1$ , see [5] and  $P$  has no infinite anti-chains see [5]. Next, by Ben Dushnik-Miller theorem, see [4], either there is an infinite set of incomparable elements in  $(P, \leq)$  or there is a chain of size  $\aleph_1$  in  $(P, \leq)$ . Now since all antichains in  $(P, \leq)$  are finite, it follows that there is a chain  $C$  in  $(P, \leq)$  of size  $\aleph_1$ . Thus, since  $C$  is scattered,  $\omega_1$  or  $\omega_1^*$  embeds in  $(C, \leq)$  see [7]. Therefore there are at least two limit points in  $\text{Fs}(P)$  which is a contradiction since the set of limit points of  $\mathcal{I}d(T \cup \{\infty\})$  is reduced to  $\{\infty\}$ .  $\square$

**Acknowledgements.** The authors thank the referee for his suggestions to use final segments instead of increasing functions as well as his fruitful remarks.

## References

- [1] U. Abraham, R. Bonnet, W. Kubis, and H. SiKaddour, *On poset Boolean algebras* (to appear in Order).
- [2] M. Bekkali, *On superatomic Boolean algebras*, Boulder, Colorado, U.S.A., 1991, Ph.D. dissertation.
- [3] M. Bekkali and D. Zhani, *Stone spaces and Ideals of posets*, International Conference (JIM' 2003), Discovery and Discrete Mathematics, Metz University, France, pp. 131–136, ISBN 2-7261-1256-0.
- [4] B. Dushnik and E. W. Miller, *Partially ordered sets*, Amer. J. Math. **63** (1941), 600–610.
- [5] R. Fraïssé, *Theory of relations*, Studies in Logic and the Foundations of Mathematics, vol. 145, North-Holland Publishing Co., Amsterdam, 2000, ISBN 0-444-50542-3.
- [6] Sabine Koppelberg, *Handbook of Boolean algebras. Vol. 1*, North-Holland Publishing Co., Amsterdam, 1989, ISBN 0-444-70261-X.
- [7] J. G. Rosenstein, *Linear orderings*, Pure and Applied Mathematics, vol. 98, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1982, ISBN 0-12-597680-1.
- [8] H. SiKaddour, *Ensembles de générateurs d'une algèbre de Boole*, Université Claude-Bernard (Lyon 1), 1988, Diplôme de Doctorat.