

BILINEAR INTERPOLATION IN TRIANGLES BASED UPON A MANGERON THEOREM

G. M. Nielson and D. J. Mangeron (*)

Recibido: 9 mayo 1979

PRESENTADO POR EL ACADÉMICO NUMERARIO D. DARÍO MARAVALL
CASESNOVES

*Dedicated to old and new Spanish
Mathematical Achievements*

Methods of approximation, which interpolate to given values on the boundary of a triangle, are presented. These methods have linear weights and are characterized on the basis of minimizing pseudonorm. Some forthcoming research projects are finally indicated. They will be related in part with some recent Ubiratan D'Ambrosio and Dicesar Lass Fernández research works.

Se exponen métodos de interpolación basados en la minimización de pseudo-normas.

1. Introduction

The fundamental bilinear interpolant of Mangeron [2], which interpolates to f on the boundary of

$$\begin{aligned} M[f](x, y) = & (1-y)f(x, 0) + yf(x, 1) + (1-x)f(0, y) + xf(1, y) - \\ & - (1-x)(1-y)f(0, 0) - x(1-y)f(1, 0) - y(1-x)f(0, 1) - xyf(1, 1) \end{aligned} \quad (1.1)$$

has had wide spread influence in the area of approximation of

$$R = \{(x, y) : 0 \leq x \leq 1; 0 \leq y \leq 1\}$$

(*) The author wishes to express his warmest thanks to the State University of Campinas Mathematics, Statistics and Computer Science Institute for the invaluable conditions offered to him to develop his work there.

bivariate functions. While $M[f]$ and its generalization to include interpolation to normal derivatives serves well for a rectangular domain, many applications require approximations which are not inherently rectangular and consequently there is interest in developing analogous methods for other domains. Of particular interest, due to the application in finite element analysis and scattered data interpolation, is a triangular domain. In this report, we present methods for a triangular domain which are patterned after the following two equivalent properties that characterize $M[f]$.

CHARACTERIZATION 1.— $M[f]$ is the unique function in $C^{2,2}(\mathbb{R})$ which lies in the kernel of the operator $\partial^4/\partial x^2 \partial y^2$ and interpolates to f on the boundary of R .

CHARACTERIZATION 2.—Among all functions in $C^{2,2}(\mathbb{R})$ which interpolate to f on the boundary of R , $M[f]$ uniquely minimizes the pseudonorm

$$\sqrt{\int_R \int [h_{xy}(s, t)]^2 ds dt}.$$

2. Linear Interpolation for Triangles

For notational convenience, we will use the standard triangular domain T with vertices $(0, 0)$, $(0, 1)$, and $(1, 0)$. Any other triangular domain can be mapped linearly to T by the use of an affine transformation. By $C^{n,m}(T)$, we mean those functions which possess continuous partial derivatives

$$\partial^{i+j} f / \partial x^i \partial y^j; \quad i \leq n, j \leq m \text{ on } T.$$

In the development of techniques of interpolation for triangular domains, there is a tendency towards the use of rational weight functions. This is due, in part, to the fact that M can be viewed as the boolean sum of two essentially univariate operators which are based upon linear interpolation along lines parallel to the sides of R . That is

$$M = M_x \oplus M_y = M_x + M_y - M_x M_y$$

where

$$\begin{aligned} M_x[f](x, y) &= (1-x)f(0, y) + xf(1, y) \\ M_y[f](x, y) &= (1-y)f(x, 0) + yf(x, 1). \end{aligned}$$

The corresponding linear interpolation operators for the domain T have rational weight function. For example,

$$T_x[f](0, y) = [(1-x-y)/(1-y)]f(0, y) + [x/(1-y)]f(1-y, y)$$

is based upon linear interpolation along lines parallel to the x -axis and so it is clear that any scheme based upon the Boolean sum of these operators will have rational weight functions. We now proceed to develop a method which is void of rational weight functions.

Concerning the first type of characterization of $M[f]$, the general form of a function $k(x, y)$ which has the property that

$$\partial^4 k / \partial x^2 \partial y^2 = 0$$

is

$$k(x, y) = g_1(x) + g_2(y) + y g_3(x) + x g_4(y)$$

for arbitrary, twice differentiable functions

$$g_i, i = 1, 2, 3, 4.$$

If we now select,

$$g_i, i = 1, 2, 3, 4$$

by forcing k to interpolate to f on the boundary of T, we obtain the equations

$$\begin{aligned} g_1(x) + g_2(0) + x g_4(0) &= f(x, 0) \\ g_1(0) + g_2(y) + y g_3(0) &= f(0, y) \\ g_1(x) + g_3(1-x) + (1-x) g_3(x) + x g_4(1-x) &= f(x, 1-x). \end{aligned}$$

Solving these equations, leads to

$$\begin{aligned} k[f](x, y) &= \frac{2-y-2x}{2(1-x)} f(x, 0) + \frac{2-2y-x}{2(1-y)} f(0, y) + \\ &+ \frac{(x+y)(3-x-y)}{2(1-x)(1-y)} f(0, 0) + \frac{x}{2(1-y)} [f(1-y, y) - f(1-y, 0)] + \\ &+ \frac{y}{2(1-x)} [f(x, 1-x) - f(0, 1-x)] + x y [g(x) - g(1-y)] \end{aligned}$$

which interpolates to f on the boundary of T for arbitrary g . In order to eliminate the rational weight functions, we choose

$$g(x) = \frac{1-2x}{2x(1-x)} [f(x, 1-x) - f(0, 1-x) - f(x, 0) + f(0, 0)]$$

which leads to the interpolation formula

$$\begin{aligned} B[f](x, y) = & (1-y)f(x, 0) + (1-x)f(0, y) - (1-x-y)f(0, 0) + \\ & + x[f(1-y, y) - f(1-y, 0)] + y[f(x, 1-x) - f(0, 1-x)]. \end{aligned} \quad (2.2)$$

It is interesting that this choice of g not only yields an interpolant with linear weights, but it also leads to an approximation which has a property analogous to the second characterization of $M[f]$.

THEOREM 2.1.—Among all functions in $C^{2,2}(T)$ which interpolate to f on the boundary of T , $B[f]$ uniquely minimizes $\langle \cdot, \cdot \rangle$ where

$$\langle h, g \rangle = \int_T \int h_{xy}(s, t) g_{xy}(s, t) ds dt.$$

PROOF.—First let h be any function in $C^{2,2}(T)$ which is zero on the boundary of T ; then by using integration by parts we have

$$\begin{aligned} \langle h, B[f] \rangle &= \int_0^1 \int_0^{1-t} h_{xy}(s, t) \frac{\partial^2 B[f]}{\partial s \partial t}(s, t) ds dt = \\ &= \int_0^1 h_y(1-t, t) \frac{\partial^2 B[f]}{\partial s \partial t}(1-t, t) dt - \int_0^1 h_y(0, t) \frac{\partial^2 B[f]}{\partial s \partial t}(0, t) dt - \\ &\quad - \int_T h_{xy}(s, t) \frac{\partial^3 B[f]}{\partial s^2 \partial t}(s, t) ds dt \end{aligned} \quad (2.3)$$

where

$$\begin{aligned} \frac{\partial^2 B[f]}{\partial s \partial t}(s, t) = & f_x(s, 1-s) - f_y(s, 1-s) - f_x(s, 0) + f_y(0, 1-s) + \\ & + f_y(1-t, t) - f_x(1-t, t) - f_y(0, t) + f_x(1-t, 0). \end{aligned} \quad (2.4)$$

Since $h(0, t) = 0$ we have that $h_y(0, t) = 0$ and from (2.4), we can see that

$$\frac{\partial^2 B[f]}{\partial s \partial t}(1-t, t) = 0$$

and so the first two terms of the right side of (2.3) vanish. Using integration by parts again, we have

$$\langle h, B[f] \rangle = \int_0^1 \int_0^{1-s} h_{xy}(s, t) \frac{\partial^3 B[f]}{\partial s^2 \partial t}(s, t) dt ds = -$$

$$= - \int_0^1 h(s, 1-s) \frac{\partial^3 B[f]}{\partial s^2 \partial t} (s, 1-s) ds + \int_0^1 h(s, 0) \frac{\partial^3 B[f]}{\partial s^2 \partial t} (s, 0) ds + \int\int_T h(s, t) \frac{\partial^4 B[f]}{\partial s^2 \partial t^2} (s, t) dt ds.$$

The first two terms are zero because $h = 0$ on the boundary of T . The last term is zero because

$$\partial^4 B[f] / \partial s^2 \partial t^2$$

is zero on T .

Let $g \in C^{2,2}(T)$ have the property that it interpolates to f on the boundary of T . Then,

$$\langle g, g \rangle - \langle B[f], B[f] \rangle = \langle g - B[f], g - B[f] \rangle + 2 \langle g - B[f], B[f] \rangle = \langle g - B[f], g - B[f] \rangle \geq 0$$

and so we have established the minimum property of $B[f]$. In order to show uniqueness, we assume the existence of another minimizing interpolant, say \bar{g} , and consider the error $e = \bar{g} - B[f]$. Since both minimize the pseudonorm, we have that

$$\langle e, e \rangle = \langle \bar{g} - B[f], \bar{g} - B[f] \rangle = 0,$$

which implies that $e_{x,y} = 0$ on T . This, along with the fact $e = 0$ on T which concludes the argument.

COROLLARY 2.2.—The operator B is exact for any function of the form

$$g(x, y) = g_1(x) + g_2(y) + xg_3(y) + yg_3(1-x)$$

for arbitrary functions $g_i, i = 1, 2, 3$.

PROOF.—Applying B directly and using

$$\begin{aligned} g(x, 0) &= g_1(x) + g_2(0) + xg_3(0) \\ g(0, y) &= g_2(y) + g_1(0) + yg_3(1) \\ g(0, 0) &= g_1(0) + g_2(0) \\ g(1-y, y) - g(1-y, 0) &= g_2(y) + g_3(y) - g_2(0) - (1-y)g_3(0) \\ g(x, 1-x) - g(0, 1-x) &= g_1(x) + g_3(1-x) - g_1(0) - (1-x)g_3(1) \end{aligned}$$

will yield this result.

In order to analyze the error and rate of convergence of this approximation, we introduce the triangular domain T_h with vertices $(0, 0)$, $(0, h)$ and $(h, 0)$ and the corresponding operator

$$B_h[f](x, y) = B[f(x/h, y/h)] \left(\frac{x}{h}, \frac{y}{h} \right) = \frac{1}{h} [(h-y)f(x, 0) + (h-x)f(0, y) - (h-x-y)f(0, 0) + x[f(h-y, y) - f(h-y, 0)] + y[f(x, h-x) - f(0, h-x)].$$

THEOREM 2.3.—For $f \in C^{1,1}(T_h)$

$$|B_h[f](x, y) - f(x, y)| \leq \frac{h^2}{2} \|f_{xy}\|_h, \quad (x, y) \in T_h.$$

where the norm is the uniform norm on T_h .

PROOF.—Since it is true in general for $f \in C^{1,1}(T_h)$ that

$$f(x, y) - f(x, 0) - f(0, y) + f(0, 0) = \int_0^x \int_0^y f_{xy}(s, t) \, ds \, dt,$$

it follows that

$$B_h[f](x, y) = \frac{y}{h} \int_0^x \int_0^{h-x} f_{xy}(s, t) \, ds \, dt + \frac{x}{h} \int_0^{h-y} \int_0^y f_{xy}(s, t) \, ds \, dt - \int_0^x \int_0^y f_{xy}(s, t) \, ds \, dt.$$

Therefore

$$|B_h[f](x, y) - f(x, y)| \leq \frac{xy(h-x)}{h} \|f_{xy}\|_h + \frac{x(h-y)y}{h} \|f_{xy}\|_h + xy \|f_{xy}\|_h = \frac{xy(3h-x-y)}{h} \|f_{xy}\|_h.$$

This bound is maximized for $(x, y) \in T_h$ when $x = y = \frac{h}{2}$ to yield the desired result.

Our forthcoming research papers will deal with application of the triangular Mangeron interpolant to scattered data interpolation and

subsequently with trilinear interpolation in tetrahedrons based upon a Mangeron extended theorem and their applications. These papers will be related also in part with some actual Ubiratan D'Ambrosio and Dicesar Lass Fernandez research work.

References

- [1] COONS, S. A. (1967). Surfaces for Computer-Aided Design of Space Forms. Project MAC report MAC-TR-41, June.
- [2] MANGERON, D. (1932). Sopra un problema al contorno per un'equazione differenziale alle derivate parziali di quart'ordine con caratteristiche reali doppie. *Rend. Accad. Sci. Fis., Mat.*, Napoli, Ser. 4, Vol. 2, 25-40.
- [3] BIRKHOFF, G. and GORDON, W. (1968). On the draftsman and related equations. *J. Approx. Theory*, 1, 199-208.
- [4] MANGERON, D. (1973). Sur certains problèmes à la frontière polygonale non nécessairement caractéristique pour une classe d'équations aux dérivées partielles d'ordre supérieur. *Comptes rendus Acad. Sci.*, Paris, 207, 94-96, 544-546, 1022-1024.
- [5] GORDON, W. (1968). Automated Design of Free Form Surfaces. General Motors Research Series. Warren, Mich.
- [6] ESWARAN, S. (1972). A Study on certain higher Order partial differential Equations of Mangeron. Doctoral Dissertation. Dept. of Mathematics, University of Alberta, Edmonton, Alberta, Canada.
- [7] NIELSEN, G. M. (1976). A Mangeron Type Theorem for a triangular domain. SIAM. *Fall Meeting. Abstracts*. Georgia Institute of Technology, Atlanta, Georgia, pp. 18-20.
- [8] MANGERON, D. (1968), Problèmes concernant les équations polyvibrantes. *Comptes rendus Acad. Sci.*, Paris, 266A, 870-873, 976-979, 1050-1052, 1103-1106, 1121-1124.
- [9] OGUZTORELI, M. N. and LEUNG, K. V. Numerical solutions of various problems concerning polyvibrating or polywave equations of Mangeron. I, II, III. *Rend. Accad. Sci. Fis., Mat.*, Naples, Series IV, Vol. 40 (1972), 52-57; *Bull. Soc. Roy. Sci.*, Liège, 42 (1973), 269-275; *Bull. Acad. Roy. Sci.*, Belgique, Serie 5, Vol. 59 (1973), 441-447.
- [10] KRIVOSHEIN, L. E., LEUNG, K. V. and OGUZTORELI, M. N. (1974). Problèmes concernant différentes classes d'équations intégro-différentielles non linéaires de Mangeron. *Rend. Accad. Naz. Lincei*, Rome, *Cl. sci. fis., mat. e nat.*, Series 8, Vol. 56, 151-156.
- [11] LEUNG, K. V. and OGUZTORELI, M. N. (1974). Numerical solution of a mixed problem for a polywave equation of Mangeron. *Collectanea Mathematica*. Vol. XXV, Fasc. 3, 1-11. Seminario Matemática de Barcelona.

- [12] MANGERON, D. (1978). On the Theory and Applications of polywave or polyvibrating equations, I. *Memoriile Sectiilor Stiintifice ale Academiei R. S. Romania (Social Republic of Romania Academy Scientific Sections Memoirs)*, IVth Series, Vol. 1, 67 p.
- [13] D'AMBROSIO, U. (1975). On limit boundaries of generalized surfaces. *Boll. U. M. I.*, Series 4, Vol. 11, 610-615.
- [14] CUADROS, C. M. (1973). Analisis Estadistico Multivariante y Representación Canonica de Funciones Estimables. *Collectanea Mathematica*. Barcelona.
- [15] FERNÁNDEZ, D. L. (1978). Interpolation of 2_n Banach Spaces. *Studia Math.*, 65.